



RESIDENTS FORUM

[Translated article] RF-Update on the Management of Vitiligo

FR-Actualización en el tratamiento del vitíligo

E.L. Pinto-Pulido*, P. Merlo-Gómez, I. Polo-Rodríguez

Servicio de Dermatología, Hospital Universitario Príncipe de Asturias, Universidad de Alcalá, Madrid, Spain

KEYWORDS

Vitiligo;
Pigmentation disorders;
Dermatological therapy

PALABRAS CLAVE

Vitíligo;
Alteraciones de la pigmentación;
Terapéutica dermatológica

The treatment of vitiligo can be challenging, as the available options show limited efficacy and are not free from adverse effects. Recently, the International Vitiligo Working Group published its therapeutic management recommendations.^{1,2} As a starting point, they emphasize deciding on a treatment goal with the patient based on the location, activity, and impact of their disease.¹

Active disease: stabilization/repigmentation goal

Limited vitiligo

Topical calcineurin inhibitors (TCIs) are considered the first choice for lesions of limited extent, especially on the face, neck, groin, and axillae. They should be applied once or twice daily for 6 months and may be extended to 12 months or longer in case of good response.

High- or very high-potency topical corticosteroids are also recommended, particularly for extrafacial areas with limited extent or as a second option for facial areas after TCIs. Application is suggested once daily for 3–6 months in adults and 2–4 months in children. These periods can be extended if an intermittent regimen (on a 2 weeks on/2 weeks off regimen) is chosen, which reduces local adverse effects and is the preferred option for facial areas.²

Ruxolitinib—a topical JAK inhibitor—is the first drug approved by the European Medicines Agency (EMA) and the Food and Drug Administration (FDA) for vitiligo repigmentation.² In phase 3 clinical trials, 80.3–81.4%/56.8–60.8% of patients achieved F-VASI50/T-VASI50 ($\geq 50\%$ improvement in facial/total vitiligo area scoring index) at 52 weeks.³

DOI of original article:

<https://doi.org/10.1016/j.ad.2023.11.023>

* Corresponding author.

E-mail address: elucia.pinto95@gmail.com (E.L. Pinto-Pulido).

<https://doi.org/10.1016/j.ad.2024.12.014>

0001-7310/© 2024 AEDV. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article as: E.L. Pinto-Pulido, P. Merlo-Gómez and I. Polo-Rodríguez, [Translated article] RF-Update on the Management of Vitiligo, ACTAS Dermo-Sifiliográficas, <https://doi.org/10.1016/j.ad.2024.12.014>

Extensive or rapidly progressive vitiligo

For patients with extensive or rapidly progressive vitiligo, narrowband UVB phototherapy is ranked as the first-line therapy, which can be combined with topical therapy. A total of 3 weekly sessions are recommended, which should be discontinued after 3 months if no improvement has been reported or after 6 months with insufficient improvement.

In patients with rapidly progressive vitiligo resistant to other therapies, oral mini-pulses of dexamethasone (2.5–5 mg) or betamethasone (5 mg) for 2 non-consecutive days per week for a maximum of 3–6 months may be considered.²

Inactive disease

Repigmentation goal

The previously mentioned topical therapies, phototherapy, or a combination of both can be used. In cases resistant to these therapies and with no disease progression after 12 months, surgical treatment with grafts—tissue or cellular—may be considered.

Stabilization goal

If the goal is to prevent relapses, twice-weekly application of a TCI or topical corticosteroid is recommended.¹

Depigmentation goal

Finally, for resistant and extensive cases in visible areas, depigmentation of healthy skin may be considered using monobenzone (FDA-approved), other topical depigmenting agents, laser, or cryotherapy.^{1,2}

Therapeutic innovations in the pipeline

Several therapies for this disease are under study. Recently, the results of a phase 2b clinical trial with ritlecitinib (an

oral JAK3 and TEC kinase inhibitor) were published, showing favorable results regarding safety and efficacy.⁴ Other oral JAK inhibitors (upadacitinib, povorcitinib, baricitinib) have also begun phase 2 clinical trials.² New therapeutic targets under investigation (IL-15, WNT signaling) aim to address not only immunomodulation but also repigmentation.^{2,5}

Funding

None declared.

Conflicts of interest

None declared.

References

1. Van Geel N, Speeckaert R, Taïeb A, Ezzedine K, Lim HW, Pandya AG, et al. Worldwide expert recommendations for the diagnosis and management of vitiligo: position statement from the International Vitiligo Task Force. Part 1: Towards a new management algorithm. *J Eur Acad Dermatol Venereol.* 2023;37:2173–84, <http://dx.doi.org/10.1111/jdv.19451>.
2. Seneschal J, Speeckaert R, Taïeb A, Wolkerstorfer A, Passeron T, Pandya AG, et al. Worldwide expert recommendations for the diagnosis and management of vitiligo: position statement from the international Vitiligo Task Force – Part 2: Specific treatment recommendations. *J Eur Acad Dermatol Venereol.* 2023;37:2185–95, <http://dx.doi.org/10.1111/jdv.19450>.
3. Janssen JC, Mulder EEAP, van der Veldt AAM. Two phase 3, randomized controlled trials of ruxolitinib cream for vitiligo. *N Engl J Med.* 2023;388:283, <http://dx.doi.org/10.1056/NEJMc2215181>.
4. Ezzedine K, Peeva E, Yamaguchi Y, Cox LA, Banerjee A, Han G, et al. Efficacy and safety of oral ritlecitinib for the treatment of active nonsegmental vitiligo: a randomized phase 2b clinical trial. *J Am Acad Dermatol.* 2023;88:395–403, <http://dx.doi.org/10.1016/j.jaad.2022.11.005>.
5. Richmond JM, Strassner JP, Zapata L Jr, Garg M, Riding RL, Refat MA, et al. Antibody blockade of IL-15 signaling has the potential to durably reverse vitiligo. *Sci Transl Med.* 2018;10, <http://dx.doi.org/10.1126/scitranslmed.aam7710>, eam7710.