


Journal Pre-proof

Survey of Usual Clinical Practice in Patients With Chronic Spontaneous Urticaria Treated With Omalizumab Regarding the Management of Antihistamines

E. Gómez de la Fuente F. J. Ortiz de Frutos M. Bergón Sendín G. Melé Ninot



PII: S0001-7310(25)00825-7

DOI: <https://doi.org/doi:10.1016/j.ad.2025.104549>

Reference: AD 104549

To appear in: *Actas dermosifiliograficas*

Received Date: 25 February 2024

Accepted Date: 16 April 2024

Please cite this article as: de la Fuente EG, de Frutos FJO, Bergón Sendín M, Melé Ninot G, Survey of Usual Clinical Practice in Patients With Chronic Spontaneous Urticaria Treated With Omalizumab Regarding the Management of Antihistamines, *Actas dermosifiliograficas* (2025), doi: <https://doi.org/10.1016/j.ad.2025.104549>

This is a PDF of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability. This version will undergo additional copyediting, typesetting and review before it is published in its final form. As such, this version is no longer the Accepted Manuscript, but it is not yet the definitive Version of Record; we are providing this early version to give early visibility of the article. Please note that Elsevier's sharing policy for the Published Journal Article applies to this version, see: <https://www.elsevier.com/about/policies-and-standards/sharing#4-published-journal-article>. Please also note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 Published by Elsevier España, S.L.U. on behalf of AEDV.

Sección: Cartas científico clínicas

Survey of Usual Clinical Practice in Patients With Chronic Spontaneous Urticaria Treated With Omalizumab Regarding the Management of Antihistamines

E. Gómez de la Fuente¹, F. J. Ortiz de Frutos², M. Bergón Sendín³, and G. Melé Ninot⁴

1. Servicio de Dermatología. Hospital Universitario Ramón y Cajal, Madrid, Spain.
2. Servicio de Dermatología. Hospital Universitario 12 de Octubre, Madrid, Spain.
3. Servicio de Dermatología. Hospital Universitario Gregorio Marañón, Madrid, Spain.
4. Servicio de Dermatología. Hospital Universitari Sagrat Cor, Barcelona, Spain.

Corresponding author: Enrique Gómez de la Fuente

E-mail address. enrique.gomezdelafuente@salud.madrid.org

To the Editor,

Clinical practice guidelines on the management of chronic spontaneous urticaria (CSU) include a series of recommendations that are not mandatory and depend on various circumstances related to both the physician and the patient. In addition, these guidelines leave several open questions, particularly regarding the management of antihistamines¹, which is why we sought to determine the usual clinical practice related to antihistamine management in patients with CSU, with special emphasis on those who initiate treatment with omalizumab.

Therefore, we created a 16-item questionnaire, in which only 1 response could be selected per question (except for Item# #13, which allowed multiple selections, and Item# 16, which invited open comments). The questionnaire was distributed among dermatologists and allergists across Spain with experience in managing omalizumab in CSU and was completed online fully anonymously (Table 1). A total of 73 specialists responded (77% dermatologists and 23% allergists) from different regions across Spain.

More than 80% of respondents reported intensifying antihistamine doses ($\times 3$ or $\times 4$) before initiating omalizumab treatment, as recommended by the European clinical practice guidelines¹. A total of 52% proceeded to the biologic without first trying a 2nd antihistamine, whereas 48% attempted a 2nd antihistamine before omalizumab administration. The latter strategy is recommended in certain guidelines, such as the Spanish guidelines from 2015² and the current British guidelines³, whereas the European guidelines are less explicit and allow transition to omalizumab after failure of the first antihistamine¹.

Clinical behavior becomes more heterogeneous once omalizumab is started. A substantial proportion of specialists report continuing antihistamines at various doses, although the general trend is to taper or discontinue them, particularly when the patient is well controlled or has already self-reduced or stopped antihistamines due to good response to omalizumab. Even at the initial omalizumab visit, more than half of specialists advise patients that they may taper or even discontinue antihistamines if their urticaria is well controlled. In the same

line, when omalizumab is optimized or discontinued, clinicians in this study reported decreasing antihistamine doses or discontinuing them in 33% and 38% of cases, respectively.

Nevertheless, a widespread practice is the use of antihistamines on demand, allowing patients to discontinue them if they have done so due to good control with omalizumab. This aligns with the belief of 86% of respondents that patients independently reduce or stop antihistamines when they feel well controlled, even if not explicitly instructed to do so. This belief has been supported by 2 recent studies in Spain. In one study based on patient surveys, 24% reported having independently discontinued antihistamines (71% due to improvement) after starting omalizumab⁴. In the other study, the quantity of antihistamines dispensed at pharmacies was evaluated and found to be significantly lower than the amount prescribed by their dermatologists⁵.

Finally, although there are no clearly established guidelines for omalizumab use in inducible chronic urticaria, clinical practice in 98% of cases mirrors that used for CSU.

In conclusion, although the behavior of Spanish specialists after initiating omalizumab is highly heterogeneous, the tendency is to taper or discontinue antihistamines. This likely reflects the absence of clear guideline recommendations and the lack of consistent positions in the literature. Some authors advocate maintaining antihistamine use^{6,7}, while others recommend reducing or discontinuing them, as they have previously failed and omalizumab alone is capable of controlling symptoms in most patients⁸. Still others support patient-directed, on-demand antihistamine use depending on disease control⁹, a practice that real-world data have shown to be common among a considerable proportion of patients⁴⁵.

To our knowledge, no similar study has been published. Although this survey may contain selection and recall biases, it reflects usual clinical practice among physicians who manage these patients. We did not observe differences based on geographic region or between dermatologists and allergists. Understanding real-world clinical practice may contribute to improved patient management. Further work is needed to refine antihistamine management in these patients and to establish evidence-based recommendations that can help standardize practices with the patient's benefit in mind.

Los autores declaran haber participado en la elaboración del manuscrito

Los autores declaramos no tener conflicto de interés

Acknowledgments

We thank the members of the *Grupo de Urticaria de la Sección Centro (UCEN)*, the *Xarxa d'Urticària Catalana i Balear*, the *Grupo Español en Investigación de Alergia Cutánea (GEIDAC)*, and other specialists from various regions of the country, who anonymously and altruistically contributed to this work.

Ética de la publicación

1. ¿Su trabajo ha comportado experimentación en animales?:

No

2. ¿En su trabajo intervienen pacientes o sujetos humanos?:

No

3. ¿Su trabajo incluye un ensayo clínico?:

No

4. ¿Todos los datos mostrados en las figuras y tablas incluidas en el manuscrito se recogen en el apartado de resultados y las conclusiones?:

Sí

References

1. Zuberbier T, Abdul Latiff AH, Abuzakouk M, Aquilina S, Asero R, Baker D, et al. The international EAACI/GA²LEN/EuroGuiDerm/APAAACI guideline for the definition, classification, diagnosis, and management of urticaria. *Allergy*. 2022;77:734-766. doi: 10.1111/all.15090.
2. Ferrer M, Bartra J, Giménez-Arnau A, Jauregui I, Labrador-Horrillo M, Ortiz de Frutos J, Silvestre JF, Sastre J, Velasco M, Valero A. Management of urticaria: not

- too complicated, not too simple. *Clin Exp Allergy*. 2015;45:731-43. doi: 10.1111/cea.12465.
3. Sabroe RA, Lawlor F, Grattan CEH, Ardern-Jones MR, Bewley A, Campbell L, et al. British Association of Dermatologists guidelines for the management of people with chronic urticaria 2021. *Br J Dermatol*. 2022;186:398-413. doi: 10.1111/bjd.20892.
 4. Melé-Ninot G, Serra-Baldrich E, Spertino J, Guilarte M, Ribó González P, Leonart-Bellfill R, et al. Are antihistamines still used during omalizumab treatment for chronic spontaneous urticaria? *Eur J Dermatol*. 2022; 1;32:629-631. doi: 10.1684/ejd.2022.4334.
 5. Menéndez Valdes M, Pérez Fernández E, López Estebaranz JL, Gómez de la Fuente E. El empleo de antihistamínicos durante el tratamiento con omalizumab en la urticaria crónica en práctica clínica real. Estudio observacional de 48 pacientes. *Actas Dermosifiliograf* 2024 (en prensa) <https://doi.org/doi:10.1016/j.ad.2023.08.015>.
 6. Madsen H, Mortz CG, Bindslev-Jensen C, Reilev M, Hallas J, Henriksen DP. High-dose non-sedating antihistamines are used insufficiently in chronic urticaria patients treated with omalizumab. *Clin Transl Allergy*. 2021; 11:e12085. doi: 10.1002/clt2.12085. Erratum in: *Clin Transl Allergy*. 2022 Feb 08;12(2):e12120.
 7. Lapeere H, Baeck M, Stockman A, Sabato V, Grosber M, Moutschen M et al. A retrospective analysis omalizumab treatment patterns in patients with chronic spontaneous urticaria: a real-world study in Belgium. *J Eur Acad Dermatol Venereol*. 2020; 34:127-134. doi: 10.1111/jdv.15684.
 8. Ensina LF, Arruda LK, Campos RA, Criado RF, Rodrigues Valle S, Melo JM, et al. H1-Antihistamines may no longer be necessary for patients with refractory chronic spontaneous urticaria after initiation of omalizumab. *J Investig Allergol Clin Immunol*. 2020; 24;30:145-147. doi: 10.18176/jiaci.0464.

9. Türk M, Yılmaz İ, Nazik Bahçecioğlu S, Can P, Ertaş R, Kartal D, et al. Effectiveness of as-needed antihistamines in chronic spontaneous urticaria patients under omalizumab treatment. *Dermatol Ther.* 2021;34:e14543. doi: 10.1111/dth.14543.

Table 1. Summary of Survey Responses on Usual Clinical Practice in Patients with Chronic Spontaneous Urticaria Treated With Omalizumab Regarding Antihistamine Management

Question / Response Options	%
1. Regarding the antihistamines used to treat CSU, which is your first-choice antihistamine?	
Bilastine	38.4
Cetirizine	26
Desloratadine	9.6
Ebastine	9.6
Levocetirizine	8.2
Rupatadine	5.5
Fexofenadine	2.7
2. Before switching from a first antihistamine to another treatment (another antiH or omalizumab) in inadequately controlled CSU (aimed at evaluating dose escalation):	
I directly switch if no response to standard dose	0
I double the dose	67
I triple the dose	18
I quadruple the dose	15
3. In uncontrolled CSU despite intensified antihistamines, prior to switching to omalizumab (to determine use of a 2nd antiH):	
I try a 2 nd -generation antihistamine at standard dose	1.4
I try a 2 nd antiH at intensified dose (×2)	6.8
I try a 2 nd antiH at intensified dose (×3 or ×4)	39.7
If no response to first antihistamine, go directly to omalizumab	52.1
4. How long do you wait before making any changes (dose or drug) in a non-responder to antiH?	
1–2 weeks	11
3–4 weeks	57.5
5–6 weeks	24.7
Other	6.8
5. At the initiation visit for omalizumab (usual behavior):	
I advise continuing intensified antihistamine dosing until first follow-up	45.2
I advise patients they may reduce antihistamine dose if urticaria is controlled	46.6
I advise patients they may stop antihistamines if completely controlled	8.2
Other	0
6. At follow-up after omalizumab if CSU is controlled (UAS7 ≤ 6 and/or UCT ≥ 12):	
I continue intensified antihistamine dosing	6.8
I de-intensify antihistamines (×2 or ×3)	56.2
I prescribe standard-dose antihistamines	15.1

Question / Response Options	%
I maintain current antihistamine dose regardless of level	15.1
I discontinue antihistamines and continue omalizumab alone	6.8
7. At follow-up after omalizumab if CSU is completely controlled (UAS7 = 0 and/or UCT = 16):	
I continue intensified antihistamine dosing	5.6
I de-intensify antihistamines (×2 or ×3)	27.8
I prescribe standard-dose antihistamines	9.7
I maintain current antihistamine dose	5.6
I discontinue antihistamines and continue omalizumab alone	51.4
8. In a patient on omalizumab + antiH in whom omalizumab is being de-intensified due to good response:	
I continue intensified antihistamines	5.6
I continue standard-dose antihistamines	26.4
I maintain current antihistamine dose	34.7
I discontinue antihistamines	33.3
9. In a patient on omalizumab + antiH in whom omalizumab is discontinued due to good response:	
I continue intensified antihistamines	5.5
I continue standard-dose antihistamines	23.3
I maintain current antihistamine dose	32.9
I discontinue antihistamines	38.4
10. In a well- or fully-controlled patient on omalizumab who has discontinued antihistamines on their own:	
I maintain omalizumab with antihistamines stopped	91
Other	9
11. If omalizumab must be intensified due to poor response:	
I intensify omalizumab + intensified antihistamines	60
I intensify omalizumab regardless of antihistamine dose	40
12. Do you act the same way in inducible urticarias as in CSU?	
Usually yes	98
Usually no	2
13. What behavior do you think your patients usually have regarding your instructions about antihistamines while on omalizumab? (multiple choice)	
I think they always follow my instructions	6.8
I think they follow them only when poorly controlled	28.8
I think they reduce the dose on their own when well controlled	86.3
I think they reduce the dose on their own regardless of control	9.6
14. What is your medical specialty?	
Dermatology	77
Allergology	23