

Table 2 Clinical and Epidemiological Characteristics.

	Autochthonous Population	Immigrant Population
Age at time of visit (most frequent 10-year age range)	19-65 years (20-30 and 30-40 years old)	19-53 years (20-30 and 30-40 years old)
Number of lesions	Single: 29 patients	Single: 3 patients
Duration of lesions	1 month to 7 years (6 patients more than 1 year)	1 month to 2 years (5 patients more than 1 year)
Risky sexual practices	28 patients (partner with condyloma, 7 cases)	5 patients (partner with condyloma, 1 case)
Associated STD	HIV+ status in 1 patient Gonococcal urethritis, 1 patient Latent syphilis, 1 patient Active hepatitis B, 1 patient	Gonococcal urethritis, 1 patient Residual syphilis, 2 patients Previous hepatitis B, 2 patients

Abbreviations: HIV, human immunodeficiency virus; STD, sexually transmitted disease.

References

- Jansâ JM, García de Olalla P. Salud e inmigración: nuevas realidades y nuevos retos. *Gac Sanit.* 2004;18 Supl 1:207-13.
- De Eusebio Murillo E, Zambrano Centeno B, Jaén Ola-solo P. Patología dermatológica en inmigrantes. *SEMERGEN.* 2004;30:345-55.
- Postigo C. Enfermedades de transmisión sexual e inmigración en España. *Actas Dermosifiliogr.* 2007;98:513-7.
- Porta N, San Juan J, Simai E, Baldellou R, Ara M, Zubiri ML. Análisis de la demanda asistencial de dermatología en la población inmigrante del área de salud del Hospital Miguel Servet de Zaragoza. *Actas Dermosifiliogr.* 2008;99:127-33.
- Albares Tendero MP, Belinchón Romero I, Ramos Rincón JM, Sánchez Payá J, Lucas Costa A, Pérez Crespo M, et al. Dermatoses en Latin American immigrants seen in a tertiary hospital. *Eur J Dermatol.* 2009;19:157-62.
- Taberner R, Nadal C, Llambrich A, Vila A, Torné I. Motivos de consulta dermatológica en la población inmigrante y española

- del área de salud del Hospital Son Llâtzer (Mallorca). *Actas Dermosifiliogr.* 2010;101:323-9.
- Albares MP, Belinchón I, Ramos JM, Sánchez Payá J, Betl-loch I. Estudio epidemiológico de la patología cutánea en la población inmigrante de Alicante. *Actas Dermosifiliogr.* 2012;103: 214-22.

R.M. Díaz Díaz,^{a,*} C. Garrido Gutiérrez,^a
A. Navarro Cantero,^b E. Ruiz Bravo-Burguillos^b

^a Sección de Dermatología Hospital Universitario Infanta Sofía, San Sebastián de los Reyes, Madrid, Spain

^b Sección de Anatomía Patológica, Hospital Universitario Infanta Sofía, San Sebastián de los Reyes, Madrid, Spain

*Corresponding author.

E-mail addresses: rmaria.diaz@salud.madrid.org, roadiatzdiaz@terra.es (R.M. Díaz Díaz).

Unresectable Angiosarcoma Treated With Bevacizumab and Paclitaxel[☆]

Angiosarcoma irreseccable tratado con bevacizumab y paclitaxel

To the Editor:

Angiosarcoma is an aggressive tumor of endothelial origin that is associated with a poor prognosis. The effectiveness of new treatments such as bevacizumab is based on their ability to inhibit angiogenesis; specifically, bevacizumab acts on vascular endothelial growth factor (VEGF), which has been implicated in the pathogenesis of angiosarcoma.^{1,2}



[☆] Please cite this article as: Nespereira-Jato MV, Peña-Panabad C, Quindós-Varela M, García-Silva J. Angiosarcoma irreseccable tratado con bevacizumab y paclitaxel. *Actas Dermosifiliogr.* 2014;105:520-522.



Figure 1 Erythematous violaceous tumor with a 4-cm diameter and peripheral macular lesions.



Figure 2 Photograph of lesion after completion of treatment with bevacizumab and paclitaxel.

We present the case of a 68-year-old woman who consulted for an erythematous violaceous tumor with a diameter of 1 cm on the scalp. Skin biopsy was consistent with a poorly differentiated angiosarcoma. The lesion grew rapidly and within a month it had progressed to a multilobulated, ulcerated tumor affecting the left side of the scalp and the right occipital region; also present in this region were red wine-colored macules measuring 1 to 2 cm in diameter (Fig. 1).

A computed tomography (CT) scan ruled out distant disease and the patient was diagnosed with poorly differentiated angiosarcoma (T2b, N0, M0), classified as stage III according to the 2010 American Joint Committee on Cancer staging system.

Immediate surgical treatment was ruled out due to the size of the lesion, which affected over half of the scalp. The patient was treated with neoadjuvant paclitaxel (80 mg/m²/wk) and bevacizumab (10 mg/kg every 2 weeks) and the lesion responded rapidly (Fig. 2). The treatment was completed after 14 cycles (each consisting of 1 bevacizumab infusion and 2 paclitaxel infusions) and was well

tolerated. The remaining lesion was surgically removed, and no evidence of angiosarcoma was seen on histology. Despite the negative biopsy results for the area of the tumor, the patient was administered adjuvant radiation therapy (daily fractions of 2 Gy, up to a total dose of 50 Gy) combined with bevacizumab (5 mg/kg every 2 weeks) due to the persistence of a red wine-colored macule in the left preauricular region. Complete response continued following completion of the 2 treatments.

Two months later, however, the patient's general health deteriorated and clinical recurrence of the angiosarcoma was noted in the left preauricular region. The CT scan showed bone metastasis. Combined treatment with paclitaxel and bevacizumab was reinitiated, but the disease progressed and the patient died 14 months after diagnosis.

Angiosarcoma is a rare but very aggressive tumor with high metastatic potential. There are 3 forms, but angiosarcoma of the head and neck—the classic form—is the most common.³ The treatment of choice is surgical excision with wide margins followed by adjuvant high-dose wide-field radiotherapy (> 50 Gy).^{1,4,5} Chemotherapy, however, is the only option for patients with advanced or unresectable angiosarcoma, but the effectiveness of different regimens has not been demonstrated in clinical trials.^{1,2} Additionally, chemotherapy is often limited because of its toxicity.¹ The most widely used regimens are anthracyclines and taxanes, but response is limited. Prospective studies of paclitaxel in the treatment of cutaneous angiosarcoma have shown a mean survival time of 8 months and a mean disease-free period of 4 months.²

The need for new drugs with improved effectiveness and tolerance is thus clear. Interest is currently focused on angiogenesis inhibitors.

Angiosarcoma originates from vascular endothelial cells, and several factors have been implicated in its pathogenesis. One of the most important of these is VEGF-A,^{1,2} which is a growth factor involved in the development of new blood vessels through the stimulation of receptors with tyrosine-kinase activity on the surface of endothelial cells. VEGF-A overexpression by neoplastic cells, which has been demonstrated in angiosarcoma and other tumors (of the lung,

Table 1 Cases of Angiosarcoma (AS) Treated With Bevacizumab in the Literature.

Authors	Year	No. of Cases	Age, y	Diagnosis	Tumor Site	Bevacizumab Doses	Other Treatments	DFS, mo	OS, mo
Koontz et al. ⁸	2008	2	50	-AS T1N0M0	Face	5 mg/kg every 2 wk; 4 doses	RT + surgery	> 26 > 8.5	ND
			63	Epithelioid AS T2N0	Face	10 mg/kg every 2 wk; 3 doses	RT + surgery		ND
Rosen et al. ⁹	2010	1	82	AS (ND on TNM)	Face	5 mg/kg every 2 wk, 14 doses	None	7	ND
Fuller et al. ⁵	2010	1	77	AS T2bN0M0	Face and neck	ND	Paclitaxel	> 11	ND
De Yao et al. ¹⁰	2011	1	88	AS T2bN0M0	Scalp	5 mg/kg every 2 wk, 4 doses	RT	7	23
Present case	2013	1	68	AS T2bN0M0	Scalp	5 mg/kg every 2 wk; 14 doses	RT + surgery	11	14

Abbreviations: AS, angiosarcoma; ND, no data; DFS, disease-free survival; OS, overall survival.

breast, and colon/rectum) promotes angiogenesis and consequently leads to increased blood supply and faster tumor growth.^{2,6} Bevacizumab is a monoclonal antibody that targets VEGF-A, preventing it from binding to its receptor, and thereby inhibiting angiogenesis. Nonetheless, the potential for recurrence and the short disease-free periods observed suggest that there are other factors at play and that VEGF inhibition alone is not sufficient to stop the tumor from growing.⁷

Five cases of cutaneous angiosarcoma treated with bevacizumab, alone or in combination, have been reported to date. A favorable response was reported in all cases, with good tolerance and longer survival than that seen with other treatments (Table 1).

The results of the first phase II clinical trial on the use of bevacizumab alone to treat metastatic or locally advanced angiosarcoma were recently published by Agulnik et al.² The results are promising, as half of the patients showed stable disease with a mean time to disease progression of 26 weeks.

Considering the scarcity of effective treatments and the results reported to date for bevacizumab in isolated cases and clinical trials, we believe that this drug could be considered in cases similar to the one we have presented. There are also clear indications that bevacizumab used in combination with other drugs could improve the prognosis in patients with angiosarcoma.

References

- Young RJ, Brown NJ, Reed MW, Hughes D, Woll PJ. Angiosarcoma. Lancet Oncol. 2010;11:983–91.
- Agulnik M, Yarber JL, Okuno AH, von Mehren M, Jovanovic BD, Brockstein BE, et al. An open-label, multicenter, phase II study of bevacizumab for the treatment of angiosarcoma and epithelioid hemangioendotheliomas. Ann Oncol. 2013;24: 257–63.
- Armenot-Carbó M, Roca-Estellés MJ, Quecedo-Estébanez E, Gimeno-Carpio E. Angiosarcoma cutáneo tras radioterapia por cáncer de mama. Actas Dermosifiliogr. 2012;103:557–9.
- Hulyalkar R, Rakshit T, García-Zuazaga J. The role of radiation therapy in the management of skin cancers. Dermatol Clin. 2011;29:287–96.
- Fuller CK, Charlson JA, Dankle SK, Russell TJ. Dramatic improvement of inoperable angiosarcoma with combination paclitaxel and bevacizumab chemotherapy. J Am Acad Dermatol. 2010;63:e83–4.
- Shrod SS, Bressler LR, Tierney LA, Cuellar S, George A. Understanding and managing the possible adverse effects associated with bevacizumab. Am J Health-Syst Pharm. 2009;66:999–1013.
- Park MS, Ravi V, Araujo DM. Inhibiting the VEGF-VEGFR pathway in angiosarcoma, epithelioid hemangioendothelioma, and hemangiopericytoma/solitary fibrous tumor. Curr Opin Oncol. 2010;22:351–5.
- Koontz BF, Miles EF, Rubio MAD, Madden JF, Fisher SR, Scher RL, et al. Preoperative radiotherapy and bevacizumab for angiosarcoma of the head and neck: Two case studies. Head Neck. 2008;30:262–6.
- Ronsen A, Thimon S, Ternant D, Machet MC, Paintaud G, Machet L. Partial response to bevacizumab of an extensive cutaneous angiosarcoma of the face. Br J Dermatol. 2010;163:208–34.
- De Yao JT, Sun D, Powell AT, Rehms EH. Scalp angiosarcoma remission with bevacizumab and radiotherapy without surgery: A case report and review of the literature. Sarcoma. 2011;2011:160369, <http://dx.doi.org/10.1155/2011/160369>.

M.V. Nespereira-Jato,^{a,*} C. Peña-Panabad,^a
M. Quindós-Varela,^b J. García-Silva^a

^a Servicio de Dermatología Hospital Abente y Lago, Complejo Hospitalario Universitario, A Coruña, Spain

^b Servicio de Oncología, Hospital Abente y Lago, Complejo Hospitalario Universitario, A Coruña, Spain

* Corresponding author.

E-mail address: [\(M.V. Nespereira-Jato\).](mailto:vnespereira@hotmail.com)

Autochthonous Cutaneous Myiasis Due to *Chrysomya bezziana*☆

Miasis cutánea no importada por *Chrysomya bezziana*

Myiasis is a parasitic infestation of the tissues or organs of vertebrates (including humans) produced by the larvae of different species of fly (Diptera).^{1–3} It is classified accord-



ing to the association between the parasite and the host and is described as obligatory, facultative, or accidental. It can also be classified according to the anatomic site of infestation as cutaneous, intestinal, or cavitary myiasis.^{1–3} Clinically, cutaneous myiasis is divided into furuncular, migratory, and wound forms.^{1–3} Incidence is higher in tropical countries with a humid climate and a low socioeconomic level.^{1–3}

We present a case of autochthonous cutaneous myiasis caused by *Chrysomya bezziana*, a species that is exceptional in Europe.

The patient was a 56-year-old man with a personal history of obesity, occasional alcohol consumption, arterial hypertension, sleep apnea-hypopnea syndrome, and extrinsic asthma. He was a heterozygous carrier of the PT20210A mutation, although he had no history of thrombosis. He had a vascular ulcer on the left leg that first appeared 10 months

☆ Please cite this article as: Aguado Lobo M, Hernández-Núñez A, Isabel García-Arata M, Borbujo J. Miasis cutánea no importada por *Chrysomya bezziana*. Actas Dermosifiliogr. 2014;105:522–524.