

- cence imaging system. Book of Abstract of the Annual Meeting of the European Society for Photodynamic Therapy. Italy, Brescia; 2005. p. 57.
6. Olsen EA, Abernethy L, Kulp-Shorten C, Callen JP, Glazer SD, Huntley A, et al. A double-blind, vehicle-controlled study evaluating masoprocol cream in the treatment of actinic keratoses of hand and neck. *J Am Acad Dermatol.* 1991;5:738-43.
  7. Fernández-Guarino M, García-Morales I, Harto A, Montull C, Pérez-García B, Jaén P. Terapia fotodinámica: nuevas indicaciones. *Actas Dermosifiliogr.* 2007;98:377-95.
  8. Tarstedt M, Rosdahl I, Berne B, Svanberg K, Wennberg AM. A randomized multicenter study to compare two treatment regimens of topical methyl aminolevulinat (Metvix®) in actinic keratosis on the face and scalp. *Acta Derm Venereol.* 2005;85:424-8.
  9. Szeimies RM, Barrer S, Radakovic-Fijan S, Tanew A, Calzavara-Pinton PG, Zane C, et al. Photodynamic therapy using topical methyl 5-aminolaevulinat compared with cryotherapy for actinic keratosis: a prospective, randomized study. *J Am Acad Dermatol.* 2002;47:258-62.
  10. Pariser DM, Lowe NJ, Stewar DM, Jarratt MT, Lucky AW, Pariser RJ, et al. Photodynamic therapy with topical methyl aminolevulinat (Metvix®) is effective and safe in the treatment of actinic keratosis: results of a prospective randomized trial. *J Am Acad Dermatol.* 2003;48:227-32.
  11. Morton C, Campbell S, Gupta G, Keohane S, Lear J, Zaki I, et al. Intraindividual right-left comparison of topical methyl-aminolevulinat-photodynamic therapy (MAL-PDT) and cryotherapy in subject with actinic keratosis: a multicentre, randomized controlled study. *Br J Dermatol.* 2006;155:1029-36.
  12. Freeman M, Vinciullo C, Francis D, Spelman L, Nguyen R, Fergin P, et al. A comparison of photodynamic therapy using topical methyl aminolaevulinat (Metvix®) with single cycle cryotherapy in patients with actinic keratosis: a prospective, randomized study. *J Dermatol Treat.* 2003;14:99-106.
  13. Ericson MB, Sandberg C, Stenquist B, Gudmundson F, Karlsson M, Ros AM, et al. Photodynamic therapy for actinic keratosis at varying fluence rates: assessment of photobleaching, pain and primary clinical outcome. *Br J Dermatol.* 2004;151:1204-12.

## FE DE ERRATAS

En el artículo titulado “*Selectividad de la fototermólisis en el tratamiento de las manchas en vino de Oporto mediante múltiples pulsos de láser de colorante pulsado*” y firmado por I. Aldanondo, P. Boixeda, M. Fernández-Lorente, A. Marquet, M. Calvo y P. Jaén (*Actas Dermosifiliogr.* 2008;99:546-54) se omitió por error el nombre de una de las autoras del trabajo, en concreto el de la Dra. E. Martín-Sáez.