


Journal Pre-proof

Secondary intention healing after Mohs micrographic surgery: an updated review of classic and novel applications, benefits and complications

Javier Gil-Lianes Ignasi Marti-Marti Daniel Morgado-Carrasco



PII: S0001-7310(24)01058-5

DOI: <https://doi.org/doi:10.1016/j.ad.2024.09.024>

Reference: AD 4219

To appear in: *Actas dermosifilograficas*

Received Date: 22 May 2024

Accepted Date: 26 September 2024

Please cite this article as: Gil-Lianes J, Marti-Marti I, Morgado-Carrasco D, Secondary intention healing after Mohs micrographic surgery: an updated review of classic and novel applications, benefits and complications, *Actas dermosifilograficas* (2025), doi: <https://doi.org/10.1016/j.ad.2024.09.024>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 Published by Elsevier España, S.L.U. on behalf of AEDV.

Sección: Dermatología Práctica**Secondary intention healing after Mohs micrographic surgery: an updated review of classic and novel applications, benefits and complications****Cierre por segunda intención después de Cirugía Micrográfica de Mohs. Revisión actualizada de sus aplicaciones clásicas y novedosas, beneficios y complicaciones**

Authors: Javier Gil-Lianes¹, Ignasi Marti-Marti¹, and Daniel Morgado-Carrasco¹

¹Servicio de Dermatología, Hospital Clínic de Barcelona, Universitat de Barcelona, Spain.

Corresponding author:

Daniel Morgado-Carrasco

E-mail address: morgadodaniel8@gmail.com

Abstract

Although secondary intention healing (SIH) is a fundamental aspect of postoperative care following Mohs micrographic surgery (MMS), it is currently underutilized. SIH constitutes a safe, cost-effective, and versatile method for wound closure. SIH offers multiple advantages, including enhanced cancer surveillance, reduced pain, and promoting esthetic outcomes, particularly not only on certain anatomical regions such as the medial canthus, antihelix, temple, or alar crease, but also for relatively small and superficial defects on the eyelids, ears, lips, and nose, including the alar region, and defects on the hands dorsal regions. Careful patient selection and thorough risk assessment are imperative to mitigate potential complications, including retraction, hyper/hypopigmented scars, or delayed healing. This comprehensive review aims to inform evidence-based decision-making on the role of SIH in MMS, synthesizing its indications, advantages, complications, wound care, and integration with other reconstructive methods.

Keywords: Mohs surgery; surgery; secondary intention healing; reconstruction; dermatology; skin cancer

Resumen

La curación por segunda intención (CSI) es un aspecto fundamental de la Cirugía Micrográfica de Mohs (CMM), pese estar actualmente infrautilizada. CSI representa un método seguro, rentable y

versátil para el cierre tras CMM. CSI tiene múltiples ventajas: mejor vigilancia de recurrencia cancerígena, reducción del dolor y resultados cosméticos favorables, particularmente en determinadas regiones anatómicas como el canto medial, antihélix, sien o pliegue alar nasal, pero también para defectos relativamente pequeños y superficiales en párpados, orejas, labios, nariz y dorso de manos. Una selección cuidadosa de los pacientes es esencial para limitar potenciales complicaciones (retracción, cicatrices hiper/hipopigmentadas o retrasos en la cicatrización. Esta revisión tiene como objetivo proveer la toma de decisiones basada en evidencia en el manejo con CSI de defectos post-CMM sintetizando sus indicaciones, ventajas, complicaciones, cuidado de heridas e integración con otros métodos reconstructivos.

Palabras claves: Cirugía Micrográfica de Mohs; cierre por segunda intención; cirugía; reconstrucción; dermatología; cáncer cutáneo

Introduction

Originally, Mohs micrographic surgery (MMS) predominantly was used a fixed tissue technique, while defects were left to heal by secondary healing intention (SIH)¹⁻⁴. The appearance of this fresh tissue technique in the 1960-70s led to a shift towards more sophisticated methods for wound closure, with SIH currently accounting for < 25% (0.8%-37.9%) of cases reported¹⁻⁴. The seminal work by Zitelli from 1983 introduced SIH as a straightforward wound management technique which was particularly praised for its excellent esthetic outcomes on certain facial sites⁵. SIH offers enhanced cancer monitoring, simplified wound care, and low-rate of complications^{6, 7}. Recent literature has reported expanded applications of post-MMS SIH in anatomical areas previously deemed suboptimal¹⁻³. This review aims to provide an update on SIH indications and advantages, focusing on anatomical considerations.

Methods

We performed a comprehensive narrative search of the literature across PubMed and Google Scholar, from inception to April 2024 using the following key words: “Mohs”, “Mohs Surgery”, “Secondary intention healing”, “Secondary intention”. Articles with a Spanish, English or German version were included and selected according to their relevance.

A potential limitation of this review is that the choice of SIH is not significantly impacted by whether the defect is due to MMS or conventional surgery. Limiting the search to MMS may have

overlooked relevant SIH data from other surgical techniques, which is acknowledged in the findings interpretation.

Indications for secondary intention healing

There are certain cases in which SIH should particularly be considered^{1, 5, 8, 9}:

1. Tumours in high-risk areas where a delayed closure is considered, especially in concave areas.
2. Tumours with aggressive features or when MMS is especially complex: post-MMS relapses or MMS > stages 3b.

SIH could also be suitable for patients with certain comorbidities including coagulation disorders, advanced age, and social or work conditions that contraindicate complex surgery.

Contraindications

The main contraindication for SIH is exposure of sensitive anatomical structures, such as vessels or nerves⁹. Special caution should be taken in cases of previous or concomitant radiotherapy^{9, 10} (higher risk of prolonged reepithelization and radionecrosis, and SIH time-to-epithelize can interfere with optimal radiotherapy schedule), inadequate postoperative care; predicted poor functional outcomes; or patients with social or occupational obligations requiring prompt reinstatement^{1, 9, 10} (**Table 1**).

Advantages and Disadvantages of secondary intention healing

The main advantage of SIH is the efficient detection of tumour recurrence (**Table 2**). Furthermore, there is no risk of certain adverse effects (seroma, suture granuloma, secondary suture failure), a lower risk of surgical site infection (SSI), and hematoma^{1, 11, 12}.

Drawbacks include prolonged healing time (increased with compromised healing process, e.g. prior radiotherapy, diabetes and mTOR inhibitors therapy), increased risk of bleeding (especially in patients on antiplatelet/anticoagulant therapy or with coagulation disorders), and risk of retraction or poor esthetic outcomes, especially if free anatomical margins are involved.

Cosmesis

In an appropriate surgical context, SIH can result in similar or better esthetic outcomes vs surgical closure^{12,13}. The most important factor is the contour of the areas involved: more favorable results on concave profiles (**Figure 1**). Secondary factors are wound size and depth (better if small and superficial), patient age, and skin colour. SIH tends to leave hypopigmented scars, which are less visible in lighter skin phenotypes^{2,3,14,15}.

Facial concavities (medial canthus and conchal bowl) heal imperceptibly, whereas convex surfaces (nasal tip and malar cheek) can heal poorly with depressed scars. Although flat areas of the cheeks, forehead, and chin heal properly, cosmesis can be unpredictable. These regions were summarized by Zitelli⁵ as NEET (concavities of the nose, eyes, ears, and temple), NOCH (convexities of the nose, oral lips, cheek, chin, and helix), and FAIR (flat areas of the forehead, antihelix of the ear, eyelids, and rest of the nose, lips, and cheeks). However, indications for SIH have since expanded to other anatomical regions^{2,6,8} (**Figure 2**).

Combination with other repairing techniques

SIH can be combined with various repairing techniques, offering versatility in wound closure, ie wounds affecting ≥ 2 cosmetic subunits (**Figure 3**). Combination with purse string or partial closures can minimize the area requiring SIH (**Figure 4**), thus reducing healing time¹⁴. In situations of uncertainty, SIH can be employed, and esthetic outcomes later be evaluated. This approach reveals new options as the wound becomes smaller and more vascularized¹⁴.

Complications

The rates of postoperative complications with SIH are low ($< 3\%$), and probably less common than with other closure techniques^{1,2,16}. SIH is associated with a comparatively lower risk of complications such as hematoma¹, patients exhibit less postoperative pain¹⁷, and SSI happen to be a rare finding (0.7% up to 4.2%)¹⁸⁻²².

Failure to re-epithelize may be due to various factors (epidermal maturation arrest, persistent granulation tissue, deficient blood supply, or infection) and wound contraction can lead to retraction and unfavorable cosmesis, particularly at free anatomical borders, ie, ectropion in the palpebral region. Other rarer potential complications include eyelid notching/webbing, trichiasis, telangiectasia, hemorrhage, bone necrosis, osteomyelitis, depressed scars, and hyperplastic granulation¹.

Antibiotic prophylaxis and topical antibiotics

Current clinical practice guidelines specify that pre- or perioperative antibiotics should be prescribed to patients who are susceptible to endocarditis and prosthetic joint infection after surgical procedures in contaminated areas, such as the oral mucosa, infected non-oral sites, or high-risk of local infection²³. A recent meta-analysis showed no statistically significant reduction in SSI in MMS after oral antibiotic prophylaxis vs placebo²⁴. Specifically in SIH, in a randomized clinical trial with 84 patients undergoing SIH on the auricular regions, no difference in SSI was seen in patients with or without levofloxacin prophylaxis (2.4% vs 2.5%)²⁵. According to the 2023 position paper of the German Society of Dermatology, there is insufficient evidence to support perioperative antibiotic prophylaxis and/or the topical application of antibiotics on wounds undergoing SIH. Routine application of antibiotic-containing ointments should be avoided to prevent sensitization, in the absence of skin barrier, and antibiotic resistance^{26,27}.

Wound care

Basic wound care can be administered by the patient, a family member, or a nurse, with scheduled visits to the dermatologist for follow-up. The wound should be kept clean to be able to apply an occlusive ointment (petrolatum or similar)^{2,9,28} and a conventional or hydrocolloid dressing. Dressing change frequency—every 2 to 7 days—depends on the amount of secretion leakage, and patients should be made aware of SSI signs^{2,9}. However, the management of wound care should be individualized based on specific factors including patient age, comorbidities (ie, diabetes, peripheral vascular disease), and location and size of the defect.

Secondary intention healing after Mohs micrographic surgery on specific anatomical areas

Ear

SIH for auricular defects after MMS has been used extensively²⁹ with good esthetic and functional outcomes. In a study on 133 patients with full-thickness auricular defects (helix, antihelix, concha, pretragal, tragal area, lobule, and posterior aspect), SIH had excellent esthetic outcomes, particularly in concave areas, even if the cartilage was removed (except for the helix, where a depression persisted). Minor cartilage exposure (< 1cm) was not a contraindication. All wounds healed in less than 10 weeks.²⁹ In a recent systematic review of the reconstruction of the auricular concha, SIH was

considered a valid option although with a higher risk of SSI³⁰. In a recent study, SIH yielded esthetic outcomes similar to full-thickness skin grafts on the helix (mean diameter, 1.7 cm up to 1.9 cm), without any differences being reported in adverse events³¹. Even if there is a higher risk of depressed scars, many patients, especially the elderly, consider slight depressions cosmetically acceptable.³¹ Recent comparative studies have shown that while split-thickness skin grafts may lead to faster healing, SIH patients experienced significantly less pain³².

Nose

A retrospective study including 96 defects on the nose (nasal tip, n = 39; alar region, n = 32; sidewall, n = 17, and dorsum, n = 8), with a mean size of 0.83 cm², revealed that diameter and depth significantly impacted scar outcome ($p < 0.001$). Nasal defects < 1cm and, which did not extend beyond the superficial fat healed well with SIH regardless of their location³³. A former study with 37 patients showed better results on concave areas—nasal ala and sidewall—and worse on the nasal tip (except if small and superficial)¹² (**Figure 5**). Regarding mean healing time, a retrospective study reported 3-4 weeks for alar or nasal tip defects of 0.5 cm up to 1.5 cm in size³⁴.

Regarding the nasal ala, in patients unable or unwilling to undergo complex nasal flaps, free-cartilage batten graft (FCBG) along with SIH can be a useful alternative^{35,36}. In a retrospective study of 129 patients who underwent FCBG with SIH good to excellent results were obtained, especially in superficial or small to intermediate-sized defects, with the cartilage closely approximating defect size, as shown in former studies^{37,38}. Healing time was estimated from 6 (small/superficial defects) up to 9 weeks (deeper/larger wounds). Only 14% of patients presented alar retraction. No hematomas or infections were reported³⁵. The authors concluded that FCBG with SIH may be considered in mid-alar wounds that are relatively shallow—>4 mm from the alar rim—and filled with a cartilage graft that is 75% up to 100% of the defect size^{39,40}.

The nasal tip does not universally heal well after SIH due to the risk of asymmetries and atrophic scars, and most surgeons prefer other surgical procedures⁴¹. SIH in the alar rim should be used with caution, especially where there are large or deep defects, since there is a risk of retraction, poor cosmesis, and collapse^{5,42}.

Lips

Classically, SIH was considered in vermilion-only and partial-thickness defects (superficial involvement of the orbicularis muscle)^{43, 44}. In a study with 68 cases of vermilion defects (mean size, 1.2 cm²) patients achieved excellent functional outcomes with good cosmesis (87% of patients would choose SIH again)¹¹, even for vermilion defects as large as 2.8 cm², or involving cutaneous lips (22/68) and/or muscular layers (23/68). A similar study with 25 patients with intermediate and large partial thickness defects (mean size, 1.6 cm), showed good to excellent esthetic and functional outcomes⁴⁴. Smaller case series revealed similar results⁴⁴⁻⁴⁶. Reported mean healing time for intermediate/large partial thickness defects on the lips was 25 days⁴⁴. SIH of the vermilion can also be combined with lateral advancement flaps if the defect involves > 2 mm of cutaneous lip^{45, 46}. Defects extending deeper than the superficial orbicularis muscle may result in esthetic or functional deformities⁴⁶ and other surgical techniques should be considered alone or in combination with SIH⁴⁵. Regarding the upper lip—with no involvement of the vermilion—a study with 105 patients with lip and chin defects showed satisfactory healing for the alar base and upper lip¹³. The apical triangle is the superior tip of the upper lip, bound by the medial cheek, nasal ala, and a hypothetical border extending from the nasolabial fold. A retrospective study (n = 24) confirmed good esthetic outcomes with SIH, with no statistically significant differences vs immediate closure⁴⁷.

Periocular region

SIH has traditionally been used for small and concave wounds, such as on the medial canthus⁴⁸. However, a retrospective study on 39 periocular wounds: lower eyelid, n = 14; upper eyelid, n = 12; lateral canthus, n = 6, and medial canthus, n = 7, and defects < 1.04 cm² showed good outcomes. Anatomic location, eyelid margin involvement and age were not significantly associated with esthetics outcomes.⁴⁹ Lowry et al⁵⁰ reported its use on 59 patients with defects ranging from 3.3 mm up to 22.3 mm on the periocular region: medial canthus, n = 32; lower eyelid, n = 20; upper eyelid, n = 4; glabella, n = 2; and nasojugal fold, n = 1. Five defects involved the eyelid margin, and 3 the canalicular system. Favorable functional and esthetic outcomes were achieved in 83% of individuals, with complications occurring in 10/59 patients: ectropion, medial canthal webbing, trichiasis, eyelid notching and hypertrophic scarring, with only 2 requiring secondary repair. Trieu et al⁵¹ reported the use of SIH on the lower eyelid in small defects (0.09 cm² up to 1.38 cm²) on 17 patients, with 100% patient satisfaction with the esthetic outcomes achieved. There was only 1 case of trichiasis, and all defects healed by week 2.

Overall, use of SIH in the periorbital region can be safe and effective, especially if the defect is < 1cm² (or < 25% of the eyelid) and superficial, regardless of location and eyelid involvement⁵¹.

Scalp and forehead

SIH represents a valid primary reconstructive option for forehead and scalp defects, especially in balding scalps⁵². Becker et al⁵³ evaluated 135 patients who had full-thickness defects on the forehead. Defects in the central area healed with atrophic, white, and depressed scars, while defects in the glabellar and temporal regions healed better. SIH can be used for large defects on the scalp (>10cm in diameter)⁵⁴. Regarding healing time, Daly et al. reported a re-epithelization time of 3 to 4 weeks for smaller wounds (< 2 cm diameter) and 6 weeks for intermediate wounds (2 cm up to 5 cm)⁵². For wounds with exposed bone, especially without periosteum, SIH may be preferable to surgical reconstruction. In such cases, fenestration of the bone cortex promotes granulation tissue and subsequent healing⁵⁵. Biosynthetic collagen dressings can also be useful⁵⁶.

In a study with 205 patients undergoing SIH after MMS on the scalp and forehead, 38 patients exhibited bone exposure with a mean area of 10.7 cm². In those cases, mean time to re-epithelialize was 13 vs 7 weeks when the periosteum was preserved. A similar retrospective study with 41 patients with defects with exposed bone on the scalp, forehead or temple showed a mean time to complete granulation of 92 days (186 days for re-epithelization). Good cosmesis was achieved in 57% of cases and no SSI were reported⁵⁷. In a study of 91 patients with exposed bone defects on the head healed by SIH, only 2.7% of patients experienced SSI, and 0% cases of osteomyelitis were observed¹⁹.

Defects on the eyebrows and above left minimal distortion, even in cases of large and deep defects. However, 4 large defects affecting contiguous subunits and/or involving muscle, periosteum, or bone caused eyebrow distortion⁵³. A smaller case series showed similar results⁵⁸, with good cosmesis, although telangiectasias were relatively common.

Cheek

Convex anatomical regions, such as the cheek, are traditionally considered not optimal for SIH. However, a study on 132 wounds on the cheek⁵⁹ (wound size from 6.3 cm² up to 32.5 cm², and depth up to subcutaneous layer, parotid gland or muscular structures in nasolabial folds) showed that most defects healed after 3 to 6 weeks. SIH in the nasolabial fold and preauricular areas achieved excellent

results⁵⁹ (**Figure 5**). Conversely, only half of the defects in the cheek medial area healed well, and defects on the mandibular or zygomatic areas healed unpredictably and often poorly¹³. Retraction tended to occur when defects extended far onto the lip or on zygomatic defects extending towards the lower eyelid⁵⁹.

Hands

A case-series of 48 patients undergoing SIH on the dorsal aspect of the hands (n = 37) or fingers (n = 11) after MMS (0.8–6 cm) showed no functional changes, with most patients reporting excellent or good cosmesis. None of the defects crossed joints or involved exposed tendons without paratenon. The authors also mention the combination of SIH plus purse string or partial closures to minimize the area left to SIH¹⁸. Another case series with 28 full-skin thickness defects involving the fascia or subcutaneous fat, with no tendon exposed and a median size of 2.4 cm (1.5 cm up to 4.6 cm), revealed a median time of healing of 44 days, and a high rate of patient satisfaction. As for AE, overgranulation developed in 12 of the 28 wounds, which resolved after applying a topical corticosteroid and discontinuing hydrocolloid dressing⁶⁰.

Lower extremities

The plantar region can be a complex site to repair. A retrospective study of 25 patients with melanoma on the soles compared 13 patients treated with SIH and 12 repaired using full-thickness skin graft. Esthetic, functional, and clinical outcomes were more favorable with SIH, although wounds took longer to heal (12 vs 8 weeks), without any differences being reported in side effects. Such findings have been previously reported^{61–63}.

Genital area

In a retrospective study on 20 patients with penile tumors treated with MMS, 80% were left to heal by SIH with good esthetic outcomes⁶⁴.

Conclusions

SIH represents a straightforward, safe, well-established, and cost-effective^{1, 2, 12} method of wound healing⁶. This approach—characterized by basic outpatient postoperative care—has a low infection

rate, preserves local skin architecture, and enables swift visualization and detection of recurrence in the management of recurrent, aggressive, and/or previously treated tumors. Several critical factors, including defect location, size, depth, geometry and colour must be meticulously considered to guarantee optimal outcomes^{1,38}. While smaller or superficial defects in concave areas often yield superior results³⁸, SIH can achieve favorable outcomes in the periocular region, lips, and nose too, including the alar region, ears and dorsal aspect of hands. Furthermore, SIH can achieve better functional and esthetic outcomes than flaps or grafts^{1,50,51}. Moreover, SIH allows potential subsequent surgical procedures or combinations with other closure techniques. The major drawbacks of SIH are the long postoperative care needed, particularly with large defects, and wound retraction, particularly at free anatomical borders, while primary contraindication remains the exposure of sensitive structures, such as nerves and arteries.

Journal Pre-proof

References

1. Moreno-Arias G, Izento-Menezes C, Carrasco M, Camps-Fresneda A. Second intention healing after Mohs micrographic surgery. *J Eur Acad Dermatol Venereol*. 2000;14:159-65.
2. Barrera-Llaurador J, Carrasquillo OY, Santiago-Vázquez M, González-Molina VJ, Valentín-Nogueras SM. Second-Intention Healing in Mohs Micrographic Surgery: A Single-Center Academic Experience. *Adv Skin Wound Care*. 2022;35:375.
3. Schwartzman G, Cartron AM, Khachemoune A. Review and reappraisal of assessment parameters of second intention healing after Mohs micrographic surgery. *Arch Dermatol Res*. 2022;314:17-23.
4. Català A, Garces J R., Alegre M, Gich I J., Puig L. Mohs micrographic surgery for basal cell carcinomas: results of a Spanish retrospective study and Kaplan–Meier survival analysis of tumour recurrence. *J Eur Acad Dermatol Venereol*. 2014;28:1363-9.
5. Zitelli JA. Wound healing by secondary intention. A cosmetic appraisal. *J Am Acad Dermatol*. 1983;9:407-15.
6. Anderson M, Flores K, Varma R. Expanded Use of Second Intention Healing in Convex Sites After Mohs Micrographic Surgery: A Focus on Cost Savings. *Dermatol Surg*. 2021;47:1123.
7. Stebbins WG, Gusev J, Higgins HW, Nelson A, Govindarajulu U, Neel V. Evaluation of patient satisfaction with second intention healing versus primary surgical closure. *J Am Acad Dermatol*. 2015;73:865-867.
8. Vedvyas C, Cummings PL, Geronemus RG, Brauer JA. Broader Practice Indications for Mohs Surgical Defect Healing by Secondary Intention: A Survey Study. *Dermatol Surg*. 2017;43:415.
9. Bechara FG. Sekundäre Wundheilung nach tumorchirurgischen Eingriffen im Kopf-Hals-Bereich. *Hautarzt*. 2013;64:567-76.
10. Sebastian G, Herrmann A. Secondary healing of the face. *Hautarzt Z Dermatol Venerol Verwandte Geb*. 2005;56:423-9.
11. Donigan JM, Millican EA. Cosmetic and Functional Outcomes of Second Intention Healing for Mohs Defects of the Lips. *Dermatol Surg*. 2019;45:26.
12. Liu KY, Silvestri B, Marquez J, Huston TL. Secondary Intention Healing After Mohs Surgical Excision as an Alternative to Surgical Repair: Evaluation of Wound Characteristics and Esthetic Outcomes. *Ann Plast Surg*. 2020;85:S28.
13. Becker GD, Levin BC, Adams LA. Outcome analysis of mohs surgery of the lip and chin: Comparing secondary intention healing and surgery. *The Laryngoscope*. 1995;105:1176-83.
14. Lam TK, Lowe C, Johnson R, Marquart JD. Secondary Intention Healing and Purse-String Closures. *Dermatol Surg*. 2015;41:S178.
15. Mott KJ, Clark DP, Stelljes LS. Regional Variation in Wound Contraction of Mohs Surgery Defects Allowed to Heal by Second Intention. *Dermatol Surg*. 2003;29:712-22.
16. Rutkowski D, Littlewood Z, Touyz SJJ, Collier NJ, Madan V, Ghura HS, et al. Prevalence of wound complications following Mohs micrographic surgery (MMS): a cross-sectional study of 1000 patients undergoing MMS and wound repair in a UK teaching hospital. *Clin Exp Dermatol*. 2022;47:1536-42.
17. Firoz BF, Goldberg LH, Arnon O, Mamelak AJ. An analysis of pain and analgesia after Mohs micrographic surgery. *J Am Acad Dermatol*. 2010;63:79-86.
18. Bosley R, Leithauser L, Turner M, Gloster HM. The Efficacy of Second-Intention Healing in the Management of Defects on the Dorsal Surface of the Hands and Fingers after Mohs Micrographic Surgery. *Dermatol Surg*. 2012;38:647-53.
19. Snow SN, Stiff MA, Bullen R, Mohs FE, Chao W-H. Second-intention healing of exposed facial-scalp bone after Mohs surgery for skin cancer: Review of ninety-one cases. *J Am Acad Dermatol*. 1994;31:450-4.

20. Ruiz-Salas V, Sanmartin-Jiménez O, Garcés JR, Vilarrasa E, Miñano-Medrano R, Escutia-Muñoz B, et al. Complications Associated with Mohs Micrographic Surgery: Data from the Nationwide Prospective Cohort REGESMOHS. *Dermatol Basel Switz.* 2022;238:320-8.
21. Schimmel J, Belcher M, Vieira C, Lawrence N, Decker A. Incidence of Surgical Site Infections in Second Intention Healing After Dermatologic Surgery. *Dermatol Surg.* 2020;46:1492.
22. Molina GE, Yu SH, Neel VA. Observations Regarding Infection Risk in Lower-Extremity Wound Healing by Second Intention. *Dermatol Surg.* 2020;46:1342.
23. Wright TI, Baddour LM, Berbari EF, Roenigk RK, Phillips PK, Jacobs MA, et al. Antibiotic prophylaxis in dermatologic surgery: Advisory statement 2008. *J Am Acad Dermatol.* 2008;59:464-73.
24. Mourad A, Gniadecki R, Taher M. Oral and Intra-incisional Antibiotic Prophylaxis in Mohs Surgery: A Systematic Review and Meta-analysis. *Dermatol Surg.* 2020;46:558.
25. Mailler-Savage EA, Neal Jr KW, Godsey T, Adams BB, Gloster Jr HM. Is Levofloxacin Necessary to Prevent Postoperative Infections of Auricular Second-Intention Wounds? *Dermatol Surg.* 2008;34:26-31.
26. Balakirski G, Becker SL, Hartmann D, Kofler L, Kunte C, Müller CSL, et al. Perioperative antibiotic prophylaxis in skin surgery – Position paper of the Antibiotic Stewardship working group of the German Society for Dermatologic Surgery (DGDC), Part 2: Special indications and situations. *JDDG J Dtsch Dermatol Ges.* 2023;21:1109-17.
27. Norman G, Dumville JC, Mohapatra DP, Owens GL, Crosbie EJ. Antibiotics and antiseptics for surgical wounds healing by secondary intention. *Cochrane Database Syst Rev.* 2016;3.
28. Yong AA, Goh CL. Use of silicone gel to enhance skin wound healing by secondary intention following tumour excision on the scalp and extremities. *Clin Exp Dermatol.* 2018;43:723-5.
29. Levin BC, Adams LA, Becker GD. Healing by Secondary Intention of Auricular Defects After Mohs Surgery. *Arch Otolaryngol Neck Surg.* 1996;122:59-66.
30. Moreno-Vazquez S, Antoñanzas J, Oteiza-Rius I, Redondo P, Salido-Vallejo R. Reconstructive Procedures of the Auricular Concha after Cutaneous Oncologic Surgery: A Systematic Review. *J Clin Med.* 2023;12:6521.
31. Hochwalt PC, Christensen KN, Cantwell SR, Hocker TL, Brewer JD, Baum CL, et al. Comparison of Full-Thickness Skin Grafts Versus Second-Intention Healing for Mohs Defects of the Helix. *Dermatol Surg.* 2015;41:69.
32. Bovenberg MS, Williams PE, Goldberg LH. Assessment of Pain, Healing Time, and Postoperative Complications in the Healing of Auricular Defects After Secondary Intent Healing Versus Split Thickness Skin Graft Placement. *Dermatol Surg.* 2024;50:35.
33. Kim DN-W, Kibbi N, Christensen SR, Leffell DJ, Suozzi KC. Factors affecting outcomes of second intent healing of nasal defects after Mohs micrographic surgery. *Arch Dermatol Res.* 2023;315:67-73.
34. Wen G, Mao D, Zhu M, Hu J, Yao X, Zhang J. Secondary Intention Healing of Nasal Ala and Tip Defects: A Simple and Valuable Treatment Option. *Clin Cosmet Investig Dermatol.* 2022;15:2165-8.
35. Kim DJ, Makdisi J, Regan C, Chen P-C, Chao E, Rotunda AM. Reconstruction of Distal Nasal Defects Using Free Cartilage Batten Grafting With Secondary Intention Healing: A Retrospective Case Series of 129 Patients. *Dermatol Surg Off Publ Am Soc Dermatol Surg Al.* 2021;47:86-93.
36. Neuhaus IM, Yu SS. Second-Intention Healing of Nasal Alar Defects. *Dermatol Surg.* 2012;38:697.
37. Ibrahimi OA, Campbell T, Youker S, Eisen DB. Nonanatomic free cartilage batten grafting with second intention healing for defects on the distal nose. *J Drugs Dermatol JDD.* 2012;11:46-50.
38. van der Eerden PA, Verdam FJ, Dennis SCR, Vuyk H. Free Cartilage Grafts and Healing by Secondary Intention. *Arch Facial Plast Surg.* 2009;11:18-23.

39. Campbell T, Eisen DB. Free Cartilage Grafts for Alar Defects Coupled with Secondary-Intention Healing. *Dermatol Surg.* 2011;37:510.
40. Rotunda AM, Cabral ES. Free Cartilage Batten Graft With Second Intention Healing to Repair a Full-Thickness Alar Wound. *Dermatol Surg.* 2014;40:1038.
41. Stigall L, Zitelli J. Reconstructing the nasal tip. *Br J Dermatol.* 2014;171:23-8.
42. Levasseur Maj JG, Mellette Jr. JR. Techniques for Reconstruction of Perialar and Perialar-Nasal Ala Combined Defects. *Dermatol Surg.* 2000;26:1019-23.
43. Faulhaber J, Géraud C, Goerdts S, Koenen W. Functional and Aesthetic Reconstruction of Full-Thickness Defects of the Lower Lip After Tumor Resection: Analysis of 59 Cases and Discussion of a Surgical Approach. *Dermatol Surg.* 2010;36:859-67.
44. Leonard AL, Hanke CW. Second intention healing for intermediate and large postsurgical defects of the lip. *J Am Acad Dermatol.* 2007;57:832-5.
45. Brahe C, Wu S, Miladi A. Mucosa Does Not Matter: Successful Secondary Intention Healing of the Lip and Vermilion Border in Mohs Surgery Patients. *Dermatol Surg.* 2021;47:692.
46. Gloster HM. The use of second-intention healing for partial-thickness Mohs defects involving the vermilion and/or mucosal surfaces of the lip. *J Am Acad Dermatol.* 2002;47:893-7.
47. Oh BH, Oh Y, Nam KA, Roh MR, Chung KY. Application of secondary intention for the restoration of the apical triangle after Mohs micrographic surgery. *J Dermatol Treat.* 2021;32:418-23.
48. Archibald LK, Gupta R, Shahwan KT, Swick M, Bakker C, Mattox AR, et al. Periorbital reconstructive techniques following Mohs micrographic surgery or excisions: a systematic review. *Arch Dermatol Res.* 2023;315:1853-61.
49. Kibbi N, Khan Y, Leffell DJ, Christensen SR, Suozzi KC. Predicting outcomes following second intent healing of periocular surgical defects. *Arch Dermatol Res.* 2021;313:483-9.
50. Lowry JC, Bartley GB, Garrity JA. The role of second-intention healing in periocular reconstruction. *Ophthal Plast Reconstr Surg.* 1997;13:174-88.
51. Trieu DN, Drosou A, White LE, Goldberg LH. Outcomes of Second Intention Healing of the Lower Eyelid Margin After Mohs Micrographic Surgery. *Dermatol Surg Off Publ Am Soc Dermatol Surg Al.* 2019;45:884-9.
52. Daly S, Gaspar Z, Francis D, Coates D, Pagliaro J. Efficacy of secondary intention healing for scalp defects—Case series from a single institution. *Australas J Dermatol.* 2023;64:522-5.
53. Becker GD, Adams LA, Levin BC. Secondary Intention Healing of Exposed Scalp and Forehead Bone after Mohs Surgery. *Otolaryngol Neck Surg.* 1999;121:751-4.
54. Barklund JS, Brown M. Second Intention Healing of a Large Surgical Defect of the Scalp. *Dermatol Surg.* 2021;47:275.
55. Barry RBM, Langtry JAA, Lawrence CM. The role of cortical bone fenestration in the management of Mohs surgical scalp wounds devoid of periosteum. *Br J Dermatol.* 2009;160:1110-2.
56. Martorell-Calatayud A, Sanz-Motilva V, Nagore E, Serra-Guillén C, Sanmartín O, Echeverría B, et al. Biosynthetic Porcine Collagen Dressings as an Adjunct or Definitive Tool for the Closure of Scalp Defects Without Periosteum. *Actas Dermo-Sifiliográficas Engl Ed.* 2012;103:887-96.
57. Wong N, Zloty D. Secondary Intention Healing Over Exposed Bone on the Scalp, Forehead, and Temple Following Mohs Micrographic Surgery. *J Cutan Med Surg.* 2022;26:274-9.
58. Deutsch BD, Becker FF. Secondary Healing of Mohs Defects of the Forehead, Temple, and Lower Eyelid. *Arch Otolaryngol Neck Surg.* 1997;123:529-34.
59. Becker GD, Adams LA, Levin BC. Spontaneous Healing of Mohs Wounds of the Cheek: A Cosmetic Assessment. *Dermatol Surg.* 1998;24:1375-82.
60. Lateo SA, Langtry JAA. A prospective case series of secondary intention healing for surgical wounds on the dorsum of the hand. *Clin Exp Dermatol.* 2013;38:606-11.

61. Audrain H, Bray A, De Berker D. Full-thickness skin grafts for lower leg defects: an effective repair option. *Dermatol Surg Off Publ Am Soc Dermatol Surg Al.* 2015;41:493-8.
62. Oganesyanyan G, Jarell AD, Srivastava M, Jiang SIB. Efficacy and complication rates of full-thickness skin graft repair of lower extremity wounds after Mohs micrographic surgery. *Dermatol Surg Off Publ Am Soc Dermatol Surg Al.* 2013;39:1334-9.
63. Schoenfeld J, Wirth P, Helm T. Mohs micrographic surgery and secondary intention healing of a plantar melanoma in-situ. *Dermatol Online J.* 2017;23.
64. Brown MD, Zachary CB, Grekin RC, Swanson NA. Penile Tumors: Their Management by Mohs Micrographic Surgery. *Dermatol Surg.* 1987;13:1163.

Journal Pre-proof

Table Legends

Table 1. Precautions on the use of SIH post-MMS

Avoid in case of	Exposure of important anatomical structures (vascular structures, nerves, tendons without paratenon, ...) Absence of vascularized structures on the defect (bone exposure without periosteum) ^a
Do not recommend in case of	Local recent radiotherapy Inability to provide adequate postoperative care Prompt return to activities of daily life required Convex surfaces ^b (nasal tip, zygomatic or mandibular area, chin) Free anatomic margins (except helix) Deep defects (deeper than subcutaneous fat) or large defects ^c Vermilion lip with > 2mm extension into cutaneous lip. Risk factors for poor wound healing (vasculopathy, smoking, malnutrition, uncontrolled diabetes, infection) High risk for endocarditis or hematogenous prosthetic infection Darker skin phenotypes (IV-V)

^a SIH can be used on exposed bone or cartilage as long as there is blood supply. This can be achieved by punching holes in the cartilage or burring holes in the bone.

^b Although convex surfaces are not the optimal regions to perform SIH, it can be considered in certain cases (ie, forehead, nasal dorsum, lip, shin, scalp) in relation to the tumor and patient features.

^c Although deep and large defects can heal well in certain locations (ie, > 4 cm on the dorsal region of hands vs >1-2 cm in high-risk facial regions), they require a long period of time to re-epithelize, and other closure techniques might be preferred initially.

Table 2. Advantages and disadvantages of SIH in MMS

Outcomes	Advantages	Disadvantages
Oncological	Early diagnosis of tumor recurrence. No disturbance of anatomic planes if tumor comes back.	
Functional	Preserves functionality in areas where movement or flexibility is crucial, ie, hands and the periocular region. No significant functional deficits reported in studies examining SIH outcomes.	Risk for ectropion or eyelid notching in the periocular region (mainly in cases of large and deep defects).
Surgical	Simplified wound management. Avoidance of complex surgery. Shorter procedural time. Lower risk of surgical site infections (vs flaps and grafts) No risk of suture dehiscence, flap necrosis, or seroma.	Longer healing times. Risk of overgranulation and delayed wound healing. Risk of bleeding (especially patients on antiplatelet/anticoagulant frugs or with coagulation disorders)

	Potential combination with other closure techniques.	
Patient	Less pain.	Psychological impact of initially open defects.
Esthetic	Good/excellent outcomes in properly selected areas.	<p>Minor imperfections are common.</p> <p>Risk of depressed scar (mainly convex, ie, cheek and chin).</p> <p>Risk of asymmetries or retraction, in areas prone to tissue movement or tension (ie, free margin borders, such as the alar rim or helix)</p> <p>Hyper- or hypopigmentation, especially in darker phototypes (IV-V)</p>

Journal Pre-proof

Figure legends




Figure 1. Medial canthus defect (2 cm x 2 cm) after MMS (A). Reconstruction of the lower region (70% of the defect) with an island flap, SIH of the upper region (30% of the defect). Esthetic outcomes after 4 weeks with minimal crusting (B).




Figure 2. Map of esthetic outcomes of SIH depending on anatomical region ^a. Green = good outcomes; Yellow = good outcomes in selected cases; Red = poor outcomes except for superficial and small wounds.

^a Front- and side-view images were generated using DALL-E by OpenAI and then modified to indicate different coloured areas.




Figure 3. Upper lip defect (1 cm x 1.5 cm) after MMS. Reconstruction of cutaneous lip with a lateral advancement flap. Vermilion defect (1 cm x 0.5 cm) was left to heal by SIHJ (A). Complete re-epithelization after 4 weeks.



Figure 4 A-C. Partial closure together with SIH. Lower eyelid defect (2.5 mm x 0.7 mm) after MMS (A). Direct closure on the lower-lateral region, SIH on the upper-medial region (1 cm x 0.6 cm) (B). Almost complete re-epithelization 10 days later. No ectropion was reported (C). D-E. Defect on the left knee (3.5 cm x 3 cm) of a 94-year-old man. Partial closure of the defect. SIH of the

central region (1.5 cm x 1.5 cm) (D). Complete re-epithelization 40 days later. Excellent functional outcomes (E).




Figure 5. SIH on folds. A-C. Lentigo maligna on the nasolabial fold. A. Delimitation of lentigo maligna prior to MMS. The nasolabial fold defect (3 cm x 4 cm) was closed using a plication of the upper and lower borders. The central defect (2 cm x 0.6 cm) was left to SIH (B). Complete re-epithelization 4 weeks later (C). **D-E. Defect on the alar groove.** Defect (1.5 cm x 0.5 cm) after MMS. B. Almost complete re-epithelization 3 weeks later. No retraction of the nasal ala seen at the follow-up.