

# Journal Pre-proof

Úlceras digitales dolorosas en paciente en hemodiálisis

F. Martins D. Paula J. Ferreira



PII: S0001-7310(25)00114-0

DOI: <https://doi.org/doi:10.1016/j.ad.2023.09.036>

Reference: AD 4284

To appear in: *Actas dermosifiliograficas*

Received Date: 4 August 2023

Accepted Date: 30 September 2023

Please cite this article as: Martins F, Paula D, Ferreira J, Úlceras digitales dolorosas en paciente en hemodiálisis, *Actas dermosifiliograficas* (2025), doi: <https://doi.org/10.1016/j.ad.2023.09.036>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 Published by Elsevier España, S.L.U. on behalf of AEDV.

## Sección: Caso para Diagnóstico

### Úlceras digitales dolorosas en paciente en hemodiálisis Managing painful digital ulcers in a patient undergoing hemodialysis

F. Martins<sup>1</sup>, D. Paula<sup>2</sup>, and J. Ferreira<sup>2</sup>

<sup>1</sup>*Department of Dermatology, Coimbra's Hospital and University Centre, Coimbra, Portugal;* and <sup>2</sup>*Department of General Surgery, Coimbra's Hospital and University Centre, Coimbra, Portugal*

Corresponding author:

Francisco Palhavã-Martins

E-mail address: francisco.paa.martins@gmail.com

#### Case presentation

A 78-year-old woman presented with a 1-month history of painful digital vesicular lesions on her left hand which had ruptured the week before forming ulcers. Additionally, she reported local itching and paresthesias, but no fever, asthenia, or anorexia. There was no past medical history of infectious processes, drug or contact allergen exposure.

The patient exhibited stage 3 hypertension and type 2 insulin-dependent diabetes mellitus. Additionally, she exhibited peripheral vascular disease and chronic kidney disease for which she underwent hemodialysis through a left brachiocephalic arteriovenous fistula. She remained on insulin on a basal-bolus regimen of carvedilol, furosemide, nifedipine, sevelamer, atorvastatin, darbepoetin, iron oxide, folic acid and calcitriol. No drug allergies were reported.

On physical examination, the patient was afebrile and hemodynamically stable. Heart and lung auscultation were normal. Periungual digital ulcers with friable erythematous bases were identified in the left hand, especially on the 2<sup>nd</sup> and 4<sup>th</sup> fingers (Figures 1 and 2). The brachiocephalic arteriovenous fistula had a palpable thrill and no local inflammatory signs were observed. Radial pulses were rhythmic but diminished bilaterally. Acrocyanosis and decreased pedal pulses were also identified, with no apparent asymmetry between fingers. Aside from acrocyanosis, no other significant skin changes in the right hand, or on the rest of the tegument were reported.

#### Diagnosis and comments:

The most suitable diagnosis in this case would be vascular access steal syndrome, an ischemic process that affects 5%-10% of patients with brachiocephalic fistulas [1,2]. In this clinical entity, collateral circulation fails to provide sufficient distal perfusion and does not compensate for the retrograde flow (or *steal*) that occurs within the arteriovenous fistula [1,2]. Therefore, patients with diabetes or atherosclerosis are at the highest risk for this condition, which is most common in women and patients aged > 60 years [1].

The diagnosis of vascular access steal syndrome is mainly clinical and 4 stages of severity can be defined. In stage I there are only subtle signs of ischemia, such as cyanosis and

decreased temperature, with few symptoms being reported, such as paresthesias. In stage II, there is pain in dialysis or exercise, which can be tolerable (II-A) or intolerable (II-B). Stage III is characterized by pain at rest and stage IV by tissue loss, which can be limited (IV-A) or irreversible (IV-B) [1].

Aside from steal syndrome, differential diagnosis in this patient, with no past medical history of infectious processes or contact allergen exposure, includes destructive arthropathy and progression of peripheral vascular disease [2,3]. However, these 2 conditions are frequently bilateral, and would not account for the presence of painful vesicular lesions solely on the left hand [2,3].

This patient—with painful periungual ulceration on the left hand but still without cutaneous necrosis—had most probably stage IV-A steal syndrome, requiring emergency treatment to prevent progression into irreversible ischemia [1,4].

The treatment of steal syndromes involves calculating the flow through the fistula, which was 1847 mL/min in this case (measured with Doppler ultrasound). When flow exceeds 1000 mL/min, the fistula is classified as high-output, and 2 surgical treatments are recommended: precision banding or revision using distal inflow (RUDI) [4,5]. While banding means simply constricting the anastomotic area, RUDI involves closing the flow through the primary anastomosis, thus creating a new fistula with a graft, and then inserting it more distally. While the former technique can lead to fistula thrombosis due to an acute decrease in flow, the latter requires good vascular patency in the terminal circulation, a condition that was not present in this patient who already exhibited advanced peripheral vascular disease [4].

Therefore, precision banding was performed. Fifteen after surgery, the patient remained asymptomatic and the ulcers had disappeared completely (Figure 3). The arteriovenous fistula remained functional.

In conclusion, painful digital ulcers in patients with peripheral vascular disease should always prompt additional investigation to exclude acute ischemia. In those with arteriovenous fistulas, the timely recognition of vascular access steal syndrome is key to successful treatment.

#### References:

- [1] Padberg FT Jr, Calligaro KD, Sidawy AN. Complications of arteriovenous hemodialysis access: recognition and management. *J Vasc Surg* 2008; 48:55S.
- [2] Tordoir JH, Dammers R, van der Sande FM. Upper extremity ischemia and hemodialysis vascular access, *Eur J Vasc Endovasc Surg*, 2004, vol. 27 (pg. 1-5)
- [3] Skeik N, Soo-Hoo SS, Porten BR, Graber J et al. Arterial Embolisms and Thrombosis in Upper Extremity Ischemia. *Vasc Endovascular Surg* 2015; 49:100.
- [4] Beathard GA, Spergel LM. Hand ischemia associated with dialysis vascular access: an individualized access flow-based approach to therapy. *Semin Dial* 2013; 26:287.
- [5] Schanzer H, Eisenberg D. Management of steal syndrome resulting from dialysis access. *Semin Vasc Surg* 2004; 17:45.

Figure 1.



Figure 2.



Figure 3.

