

- aminolevulinic acid-photodynamic therapy (MAL-PDT) and cryotherapy in subjects with actinic keratosis: a multicentre, randomized controlled study. *Br J Dermatol.* 2006;155: 1029-36.
12. Freeman M, Vinciullo C, Francis D, Spelman L, Nguyen R, Fergin P, et al. A comparison of photodynamic therapy using topical methyl aminolevulinic acid (Metvix®) with single cycle cryotherapy in patients with actinic keratosis: a prospective, randomized study. *J Dermatol Treat.* 2003;14:99-106.
 13. Ericson MB, Sandberg C, Stenquist B, Gudmundson F, Karlsson M, Ros AM, et al. Photodynamic therapy for actinic keratosis at varying fluence rates: assessment of photobleaching, pain and primary clinical outcome. *Br J Dermatol.* 2004;151:1204-12.

ERRATA

In the article entitled "*Selectivity of Photothermolysis in the Treatment of Port Wine Stains Using Multiple Pulses With a Pulsed Dye Laser*" and signed by I. Aldanondo, P. Boixeda, M. Fernández-Lorente, A. Marquet, M. Calvo, and P. Jaén (*Actas Dermosifiliogr.* 2008;99:546-54), the name of one of the authors of the article—Dr E. Martín-Sáez—was omitted.