



# ACTAS Dermo-Sifiliográficas

Full English text available at  
[www.actasdermo.org](http://www.actasdermo.org)



## ORIGINAL ARTICLE

### Analysis of Patients With *Chlamydia trachomatis* Genital Infection in an STD Clinic<sup>☆</sup>



J.B. Repiso-Jiménez,<sup>a,\*</sup> T. Fernandez-Morano,<sup>a</sup> F. Rivas-Ruiz,<sup>b</sup> M. de Troya-Martin<sup>a</sup>

<sup>a</sup> Servicio de Dermatología, Empresa Pública Hospital Costa del Sol, Marbella, Málaga, Spain

<sup>b</sup> Unidad de Investigación, Empresa Pública Hospital Costa del Sol, Marbella, Málaga, Spain

Received 12 January 2014; accepted 21 April 2014

Available online 8 September 2014

#### KEYWORDS

Sexually transmitted infections;  
*Chlamydia trachomatis*;  
Sexual behavior

#### Abstract

**Background and objectives:** *Chlamydia trachomatis* genital infection is common in our setting and early treatment can prevent complications. The aim of this study was to report on patients diagnosed with *C trachomatis* genital infection in a sexually transmitted disease (STD) clinic.

**Material and methods:** This was a descriptive, cross-sectional, observational study of patients diagnosed with *C trachomatis* infection between 2010 and 2011. We recorded demographic data and information on sexual habits, concomitant sexually transmitted infections (STIs), and various aspects of treatment.

**Results:** In total, 12.3% of the samples analyzed were positive for *C trachomatis* genital infection. Sixty-two patients (43 men) with a mean age of 31 years were studied; 75% were heterosexual and 87% had had a sexual partner in the previous 2 months. Condom use was inconsistent in 81%, 79%, and 65% of patients who practiced vaginal, oral, and anal sex, respectively. Thirteen percent of the patients had symptoms and anogenital warts were the most common associated STI. The most widely used treatment was doxycycline.

**Conclusions:** A high prevalence of genital *C trachomatis* infection was detected in our STD clinic, and the majority of cases were found in young men. We observed a high rate of asymptomatic infection in patients who do not engage in high-risk sexual behavior and who had come to the clinic for another reason. Systematic screening of *C trachomatis* infection should be implemented in STD units to enable the early treatment of patients and their recent sexual partners.

© 2014 Elsevier España, S.L.U. and AEDV. All rights reserved.

<sup>☆</sup> Please cite this article as: Repiso-Jiménez JB, Fernandez-Morano T, Rivas-Ruiz F, de Troya-Martin M. Análisis de la población con infección genital por *Chlamydia trachomatis* en una consulta de enfermedades de transmisión sexual. Actas Dermosifiliogr. 2014;105:774-779.

\* Corresponding author.

E-mail address: [boscorepiso@gmail.com](mailto:boscorepiso@gmail.com) (J.B. Repiso-Jiménez).

**PALABRAS CLAVE**  
Infecciones de  
transmisión sexual;  
Chlamydia  
trachomatis;  
Hábitos sexuales**Análisis de la población con infección genital por *Chlamydia trachomatis* en una consulta de enfermedades de transmisión sexual****Resumen**

**Introducción y objetivos:** La infección genital por *Chlamydia trachomatis* (CT) tiene una alta incidencia en nuestro medio y su tratamiento precoz puede evitar complicaciones. Describimos la población de pacientes diagnosticados en una consulta de enfermedades de transmisión sexual.

**Material y métodos:** Estudio observacional descriptivo de corte transversal de la población diagnosticada de infección por CT entre 2010 y 2011. Se recogieron datos demográficos, hábitos sexuales, asociación con otras infecciones de transmisión sexual (ITS) y diversos aspectos del tratamiento.

**Resultados:** Encontramos positividad para CT en el 12,3% de las muestras. Se incluyeron 62 pacientes con una media de edad de 31 años. De estos 43 eran hombres y el 75% era heterosexual. El 87% de los pacientes había tenido una pareja sexual en los últimos 2 meses. El uso del preservativo se consideró deficiente en el 81, 79 y 65% de los pacientes en el sexo vaginal, oral y anal respectivamente. Aparecieron síntomas en el 13% y la ITS asociada con más frecuencia fueron las verrugas anogenitales. El tratamiento más usado fue la doxiciclina.

**Conclusiones:** La detección de CT en nuestra consulta es elevada, procediendo en su mayoría las muestras de varones jóvenes. Demostramos una alta incidencia de infección genital asintomática por CT en pacientes sin prácticas sexuales de alto riesgo y que acudieron a la consulta por otros motivos. Es necesario el despistaje sistemático de CT en la consulta de ITS para tratar de forma precoz al paciente y a sus parejas recientes.

© 2014 Elsevier España, S.L.U. y AEDV. Todos los derechos reservados.

## Introduction

Genital chlamydia, which is caused by *Chlamydia trachomatis*, is the most common sexually transmitted infection (STI) in Europe. In 2009, 343 958 cases were reported in 23 member countries of the European Union, including Spain.<sup>1</sup> *C trachomatis* infection is also the main notifiable STI in our health district (unpublished data). Because most infections are mild and asymptomatic, patients do not generally seek medical attention. Spanish groups have been warning about the need for a standardized screening program for *C trachomatis* infection in STI clinics for years.<sup>2</sup> Persistent infections in women can cause pelvic inflammatory disease, infertility, and ectopic pregnancies.<sup>3</sup> Early detection of cervical infections in women have proven to be effective in reducing complications.<sup>4</sup> In men, chlamydia mainly affects the urethra and while it is associated with few complications, there is a high risk of the infection being spread to female partners.<sup>5</sup> The true prevalence of *C trachomatis* infection and associated risk factors is unknown in Spain and has been investigated in just a few studies.<sup>6</sup> The aim of this study was to determine the prevalence of *C trachomatis* infection in patients seen at an STI clinic run by the dermatology department of a public hospital in the south of Spain (Costa del Sol Occidental) and to describe the risk profile of patients diagnosed with this infection.

## Material and Methods

This was a descriptive, observational, cross-sectional study of patients diagnosed with chlamydia at an STI clinic between January 2010 and December 2011. The clinic forms part of Hospital Costa del Sol in Malaga, Spain, which serves a population of 387 036 inhabitants according to the municipal census of the Spanish National Statistics Institute. All patients seen at the clinic were evaluated and those who tested positive for urethral, cervical, or rectal *C trachomatis* at any time during follow-up were included in the study. Patients who did not consent to their data being collected were excluded. We recorded information on sexual preferences and habits using a questionnaire that is completed as part of the routine sexual history taken at the clinic. Diagnosis was established using the *artus* polymerase chain reaction kit (*C. trachomatis* Plus RG PCR Kit for use with QIAAsymphony SP/AS and Rotor-Gene Q Instrument, QIAGEN). Urethral and cervical samples were tested in men and women respectively, and rectal samples were tested in men and women with a recent history of anal sex. We also recorded the clinical characteristics of infection, the presence of other sexually transmitted diseases, and different aspects of treatment. We performed a descriptive analysis using measures of central tendency and dispersion for quantitative variables and distribution of frequencies for qualitative variables.

## Results

In total, 729 samples (388 urethral, 181 cervical, and 160 rectal) were collected from 560 patients during the study period; 90 samples (12.3%) were positive for *C trachomatis* infection in 78 patients. Of these patients, 62 (43 men and 19 women) correctly completed the questionnaire and were included in the study. Their mean age was 30.8 years (95% CI, 28.4–33.3) and most of the patients were of Spanish origin (71%) and white (82.3%). Ninety-five percent of patients had completed some level of education (lower than university level in 79.3% of cases) (Table 1). The most common professions were in the hotel, restaurant, and catering industry.

Seventy-five percent of the patients were heterosexual and 25% reported having had more than 5 sexual partners in the past year. Eighty-seven percent of those evaluated had had at least 1 sexual partner in the past 2 months, and 70% had a stable partner at the time of diagnosis. Condom use was inconsistent in 81% of those who engaged in vaginal sex, 79% of those who engaged in oral sex, and 65% of those who engaged in anal sex (Table 1).

None of the patients reported working as sex workers injecting drugs, or exchanging sex for drugs. Two men stated that they had paid for sex in the past year.

Thirteen percent of patients had chlamydia-related symptoms. Thirty-nine (90.7%) of the 43 men had a positive urethral sample and 6 of these (15.4%) had symptoms of dysuria and urethral discharge. One of the 18 women with a positive cervical sample had symptomatic cervical discharge. Genetic material of *C trachomatis* was isolated in the rectum of 4 men who have sex with men (MSM) and 3 heterosexual women, all of whom were asymptomatic. Ten (17.5%) of the 57 patients who tested positive for chlamydia and underwent serology had antibodies to *C trachomatis*; in 80% of cases, the antibodies were immunoglobulin M. None of the patients developed complications during the study period (Table 2).

Eighty-one STIs were identified in the patients studied, and 92% of patients had at least 1 other STI: 46 (80.7%) had anogenital warts, 16 (28.1%) had human papillomavirus (HPV)-related cancers, 8 (14%) had genital herpes, 5 (8.8%) had syphilis, and 2 (men) (3.5%) had urethritis caused by other pathogens. Five patients (7%) were positive for human immunodeficiency virus (HIV) infection (Table 2).

The most common treatment (used in 69.4% of cases) was doxycycline 100 mg every 12 hours for a week. Treatment was not administered in 4 patients for different reasons. Following the protocol at our clinic, all the patients were provided with information and medication for recent identifiable sexual partners. We determined that in 43 cases (69.4%), at least 1 partner received the information and was treated. Of the 32 patients who were retested after 3 months, 4 were still positive for *C trachomatis* infection (Table 3).

## Discussion

We detected a high rate of *C trachomatis* infection at our STI clinic; those infected were largely young, white adult men of Spanish origin. Our assessment of risk factors for

**Table 1** Sociodemographic Characteristics and Sexual Behavior.

	No. of Patients	% of Patients
<b>Sex</b>		
Male	43	69.4
Female	19	30.6
<b>Education<sup>a</sup></b>		
No schooling	3	5.2
Primary education	21	36.2
Secondary education	22	37.9
University education	12	20.7
<b>Geographic origin</b>		
Spain	44	71
Central and South America	12	19.4
Europe	4	6.5
North Africa	2	3.2
<b>Sexual orientation</b>		
Heterosexual	46	74.2
Homosexual	15	24.2
Bisexual	1	1.6
<b>Partners in past year, No.</b>		
< 5	47	75.8
≥ 5	15	24.2
<b>Partners in past month, No.</b>		
None	7	11.3
< 5	54	87.1
≥ 5	1	1.6
<b>Stable partner<sup>b</sup></b>		
No	18	30.5
Yes	41	69.5
<b>Anal sex</b>		
No	39	62.9
Yes	23	37.1
<b>Condom use in anal sex</b>		
Never	9	39.1
Occasional	6	26.1
Always	8	34.8
<b>Vaginal sex</b>		
No	15	24.2
Yes	47	75.8
<b>Condom use in vaginal sex</b>		
Never	11	23.4
Occasional	27	57.4
Always	9	19.2

<sup>a</sup> Information on educational level was missing for 4 patients.

<sup>b</sup> Information on stable partner was missing for 3 patients.

**Table 2** Clinical Results.

	No. of Patients	% of Patients
<b>Symptomatic</b>		
No	54	87.1
Yes	8	12.9
<b>Sample (positive)<sup>a</sup></b>		
Urethral	39	62.9
Cervical	18	29.0
Rectal	7	11.3
<b>Serology<sup>b</sup></b>		
Negative	47	82.5
Positive	10	17.5
<b>Other STI (positive)<sup>c</sup></b>		
Anogenital warts	46	80.7
HPV-related cancer	16	28.1
Genital herpes	8	14
Syphilis	5	8.8
HIV	4	7
Urethritis due to other causes	2	3.5

Abbreviations: HIV, human immunodeficiency virus; HPV, human papillomavirus; STI, sexually transmitted infection.

<sup>a</sup> For 62 patients.

<sup>b</sup> For 57 patients.

<sup>c</sup> STIs: 81 in 57 patients.

infection revealed that while a high percentage of patients were heterosexual and had just 1 sexual partner (generally a stable one) in the previous month, condom use was low. In most cases, patients were tested for *C trachomatis* infection as part of routine testing of asymptomatic patients who came to the clinic for another STI (primarily anogenital warts).

**Table 3** Treatment.

	No. of Patients	% of Patients
<b>Treatment<sup>a</sup></b>		
Doxycycline	43	69.4
Azithromycin	11	17.7
Azithromycin and doxycycline	2	3.2
None	4	6.5
<b>Partner treated</b>		
No	19	30.6
Yes	43	69.4
<b>Confirmation test</b>		
Done	32	51.6
Not done	30	48.4
<b>Result of confirmation test</b>		
Negative	28	87.5
Positive	4	12.5

<sup>a</sup> For 60 patients.

Genital *C trachomatis* infection is the main STI reported in our health district, with 238 notifications in 2 years. This is approximately twice the number of reported cases for syphilis and 5 times that for gonococcal infections. *C trachomatis* infection was the most common notifiable disease reported in the United States in 2008.<sup>7</sup> Additionally, it was detected in 7% of women aged between 15 and 24 years seen at family planning clinics between 2004 and 2008<sup>8</sup> and in 8.3% of women in the same age group in their first term of pregnancy seen at prenatal clinics.<sup>9</sup> The prevalence of chlamydia in sexually active men and women has been estimated at around 4% in some Spanish regions.<sup>6</sup> International clinical guidelines recommend testing for *C trachomatis* in sexually active women under 25 years of age,<sup>10</sup> women in their first term of pregnancy,<sup>11</sup> and men under 25 years of age.<sup>12</sup> Efficient screening programs in young women have proven successful in preventing complications due to *C trachomatis*.<sup>3,4,13</sup> Testing for chlamydia in STI clinics is justified due to the high prevalence of the infection and the fact that treatment can reduce complications.

We did not identify any particular risk groups related to race, educational level, or former residence in a developing country in our series. Studies in Europe<sup>14</sup> and the United States<sup>15</sup> have shown *C trachomatis* infection to be more common in black women, women from poor countries, and women with a lower level of education. The risk of chlamydia in young adults has been linked to inconsistent condom use<sup>6,16</sup> and an increased number of sexual partners in the preceding 3 months.<sup>16</sup> In our study, we did not find a high number of recent partners, and we assume that the infection was picked up from the patient's current partner, indicating low condom use.

In line with the findings of most studies,<sup>17</sup> the majority of patients with *C trachomatis* infection in our series were asymptomatic. The most common associated STIs in our series were HPV infections. Genital HPV infections are the main reason for referral to our clinic and do not necessarily indicate high-risk sexual behavior.

Rectal samples were positive for *C trachomatis* infection in 4 MSM and 3 women who engaged in anal sex. A high prevalence of rectal *C trachomatis* infection in patients who have had recent receptive anal intercourse has been reported in both women<sup>18</sup> and men.<sup>19</sup> Diagnosis and treatment is crucial in such cases to control the spread of infection and reduce the risk of HIV infection in MSM.<sup>20</sup>

Both doxycycline and azithromycin have proven effective in treating chlamydia, but the former is considered superior for the treatment of rectal infections.<sup>21</sup> Risk of reinfection in the year after diagnosis can be as high as 20%,<sup>22</sup> and repeat testing is therefore recommended 3 to 12 months after diagnosis.<sup>23</sup> Patient-delivered partner treatment is an effective measure for reducing risk of reinfection.<sup>24</sup> The reinfection rate in our series is similar to reports in the literature, even though appropriate treatment was delivered and partner notification was high.

While nucleic acid amplification tests have high specificity and sensitivity, bacterial culture is the most reliable method for identifying genital *C trachomatis* infection. It is, however, expensive and only available at certain reference hospitals. It should be noted that most of the patients we analyzed came to our STI clinic with another infection

(mostly genital warts), and this is an obvious source of selection bias.

We detected a high rate of *C trachomatis* infection in the population seen at our STI clinic. Considering the high prevalence and asymptomatic nature of *C trachomatis* infection, routine screening for chlamydia should be performed in all patients who consult for an STI at such clinics, regardless of their risk profile, which is generally low. We believe that routine *C trachomatis* testing should be extended to all at-risk individuals seen at health centers and gynecological clinics.

Further studies of the general population are needed to determine the true prevalence of *C trachomatis* infection. The identification of infection and treatment of both patients and their partners is key to preventing complications and transmission.

## Ethical Disclosures

**Protection of humans and animals.** The authors declare that no tests were carried out in humans or animals for the purpose of this study.

**Confidentiality of data.** The authors declare that they have followed their hospital's protocol on the publication of data concerning patients and that all patients included in the study have received sufficient information and have given their written informed consent to participate in the study.

**Right to privacy and informed consent.** The authors declare that no private patient data appear in this article.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## References

- Sexually transmitted infections in Europe 1990-2009 [consultado 18 Mar 2014]. Available in: [http://www.ecdc.europa.eu/en/publications/Publications/110526\\_SUR\\_STI\\_in\\_Europe\\_1990-2009.pdf](http://www.ecdc.europa.eu/en/publications/Publications/110526_SUR_STI_in_Europe_1990-2009.pdf)
- Cacho J, Sanz F, Blanco MA. La enfermedad silenciosa por Chlamydia trachomatis: necesidad urgente de detección y tratamiento en mujeres. Enferm Infect Microbiol Clin. 2001;19:419-21.
- Bender N, Herrmann B, Andersen B, Hocking JS, van Bergen J, Morgan J, et al. Chlamydia infection, pelvic inflammatory disease, ectopic pregnancy and infertility: Cross-national study. Sex Transm Infect. 2011;87:601-8.
- Scholes D, Stergachis A, Heidrich FE, Andriilla H, Holmes KK, Stamm WE. Prevention of pelvic inflammatory disease by screening for cervical chlamydial infections. New Engl J Med. 1996;334:1362-6.
- Pavonen J. Chlamydia trachomatis-induced urethritis in female partners of men with nongonococcal urethritis. Sex Transm Dis. 1979;6:69-71.
- Fernandez-Benítez C, Mejuto-López P, Otero-Guerra L, Margolles-Martins MJ, Suarez-Leiva P, Vazquez F, et al. Prevalence of genital Chlamydia trachomatis infection among young men and women in Spain. BMC Infectious Diseases. 2013;13:388.
- Centers for Disease Control and Prevention. Summary of notifiable diseases-United States, 2008. Morb Mortal Wkly Rep. 2008;57:1-94.
- Satterwhite CL, Grier L, Patzer R, Weinstock H, Howards PP, Kleimbaum D. Chlamydia positivity trends among women attending family planning clinics: United States 2004-2008. Sex Transm Dis. 2011;38:989-94.
- Satterwhite CL, Gray AM, Berman S, Weinstock H, Kleimbaum D, Howards PP. Chlamydia trachomatis infections among women attending prenatal clinics: United States, 2004-2009. Sex Transm Dis. 2012;39:416-20.
- US Preventive Services Task Force. Screening for chlamydial infection: US Preventive Services Task Force recommendation statement. Ann Intern Med. 2007;147:128-34.
- Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines. MMWR. 2010;59:1-110.
- Vajdic CM, Middleton M, Bowden FJ, Fairley CK, Kaldor JM. The prevalence of genital Chlamydia trachomatis in Australia 1997-2004: A systematic review. Sex Health. 2005;2:169-83.
- Oakeshott P, Kerry S, Aghaizu A, Atherton H, Hay S, Taylor-Robinson D, et al. Randomised controlled trial of screening for Chlamydia trachomatis to prevent pelvic inflammatory disease: The POPI (Prevention of Pelvic Infection) trial. BMJ. 2010;340:c1642.
- Op de Coul EL, Götz HM, van Bergen JE, Fennema JS, Hoebe CJ, Koekenbier RH, et al. Who participates in the Dutch Chlamydia screening? A study on demographic and behavioral correlates of participation and positivity. Sex Transm Dis. 2012;39:97-103.
- Torrone EA, Geisler WM, Gift TL, Weinstock HS. Chlamydia trachomatis infection among women 26 to 39 years of age in the United States, 1990 to 2010. Sex Transm Dis. 2013;40:335-7.
- Wand H, Guy R, Donovan B, McNulty A. Developing and validating a risk scoring for Chlamydia infection among sexual health clinic attendees in Australia: A simple algorithm to identify those at high risk of Chlamydia infection. BMJ Open. 2011;1:e000005, <http://dx.doi.org/10.1136/bmjopen-2010-000005>.
- Detels R, Green AM, Klausner JD, Katzenstein D, Gaydos C, Handsfield H, et al. The incidence and correlates of symptomatic and asymptomatic Chlamydia trachomatis and Neisseria gonorrhoeae infections in selected populations in five countries. Sex Transm Dis. 2011;38:503-9.
- Javanbakht M, Gorbach P, Stirland A, Chien M, Kerndt P, Guerry S. Prevalence and correlates of rectal Chlamydia and gonorrhea among female clients at sexually transmitted disease clinics. Sex Transm Dis. 2012;39:917-22.
- Wilkinson A, El-Hayek C, Fairley CK, Leslie D, Roth N, Tee BK, et al. Incidence and risk factors associated with Chlamydia in men who have sex with men: A cohort analysis of Victorian Primary Care Network for sentinel surveillance data. Sex Transm Infect. 2012;88:319-24.
- Kent CK, Chaw JK, Wong W, Liska S, Gibson S, Hubbard G, et al. Prevalence of rectal, urethral, and pharyngeal chlamydia and gonorrhea detected in 2 clinical settings among men who have sex with men: San Francisco, California, 2003. Clin Infect Dis. 2005;41:67-74.
- Hathorn E, Opie C, Goold P. What is the appropriate treatment for the management of rectal Chlamydia trachomatis in men and women? Sex Transm Infect. 2012;88:352-4.
- Hosenfeld CB, Workowski KA, Berman S, Zaidi A, Dyson J, Mosure D, et al. Repeat infection with Chlamydia and gonorrhea among

- females: A systematic review of the literature. *Sex Transm Dis.* 2009;36:478–89.
23. Centers for Disease Control and Prevention. Recommendations for the prevention and management of *Chlamydia trachomatis* infections, 1993. *MMWR.* 1993;42 (No.RR-12).
24. Trelle S, Shang A, Nartey L, Cassell JA, Low N. Improved effectiveness of partner notification for patients with sexually transmitted infections: Systematic review. *BMJ.* 2007; 334:54.