




ACTAS Dermo-Sifiliográficas

Full English text available at
www.actasdermo.org



RESIDENTS FORUM

[Translated article] RF-Adverse Skin Reaction to Apalutamide: An Emerging Effect

FR-Toxicodermia por apalutamida: una entidad emergente

M. Pons Benavent^{a,*}, S. Porcar Saura^a, L. Bou Boluda^b

^a Servicio de Dermatología, Hospital Clínico Universitario de Valencia, Blasco Ibáñez 17, Valencia 46010, Spain

^b Servicio de Dermatología, Hospital Arnau de Vilanova, San Clemente 12, Valencia 46015, Spain

KEYWORDS

Apalutamide;
Drug-related side effects and adverse reactions;
Drug eruptions;
Androgen receptor blockers

PALABRAS CLAVE

Apalutamida;
Toxicidad cutánea;
Reacción farmacológica;
Inhibidor de los receptores androgénicos

Introduction

Apalutamide is an androgen receptor inhibitor approved by the US Food and Drug Administration in 2018 for use against

certain prostate cancers. It was approved for use in Europe in March 2021.¹ Skin rashes, hypothyroidism, and bone fractures are among apalutamide's common adverse effects.¹

Case Descriptions

An 89-year-old man with a history of prostate cancer in treatment with apalutamide for 2 months complained of a skin rash that had first appeared 30 days earlier. The macules and intense peeling started on his chest and spread to involve 80% of his body (Fig. 1A). Another 89-year-old man with a history of prostate cancer also consulted us for erythematous papules over large areas of his trunk and limbs. This patient had been in treatment with apalutamide for 1 month and the lesions had appeared 7 days before he consulted us. The rash started on his chest and arms and spread upward and downward, sparing areas not exposed to the sun (Fig. 1B). Both patients reported intense itching and general discomfort. The first man was admitted to hospital because his overall condition had declined. In both cases, the biopsies demonstrated a lymphocytic infiltrate in the superficial dermis with significant damage at the dermal-epidermal junction and an abundance of apoptotic keratinocytes, suggesting an adverse drug reaction (Fig. 1C and 1D). The rashes resolved after apalutamide was stopped and a short course of systemic corticosteroids was prescribed.

DOI of original article:

<https://doi.org/10.1016/j.ad.2021.10.021>

* Corresponding author.

E-mail address: martiponsbenavent@gmail.com

(M. Pons Benavent).

<https://doi.org/10.1016/j.ad.2021.10.028>

0001-7310/© 2022 AEDV. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Figure 1 A, Erythroderma on the anterior surface of the trunk of the first patient. B, Erythematous maculopapular rash on the front of the trunk and sides of the shoulders and arms of the second patient. C and D, Skin biopsies of the chest of each patient showed a predominantly lymphocytic infiltrate in the superficial dermis, associated with significant damage at the dermal–epidermal interface and an abundance of apoptotic keratinocytes (hematoxylin–eosin, magnification $\times 200$).

Discussion

Reports of adverse skin reactions to apalutamide have been scarce in the literature to date. Such reactions are described in the summary of product characteristics for apalutamide and in a report of a series of 68 Japanese patients in clinical trials prior to approval; the adverse skin effects were described ambiguously as rashes, maculopapular rashes, and generalized rashes.^{1,2} Such skin lesions developed at one time or another in 51.7% of the patients on apalutamide, and there was a significant correlation between the presence of rashes and drug plasma concentration. In contrast, no correlation was detected between exposure indicated by plasma concentrations and symptom severity, nor between the incidence of lesions and any patient characteristics at baseline.² The frequency and severity of rashes in the Japanese series was greater than has been observed in patients in other geographic areas. A case of epidermal necrolysis due to apalutamide has been reported.³

Other androgen receptor inhibitors, however, have more often been reported to cause adverse skin reactions. Noteworthy are photosensitivity reactions associated with flutamide and bicalutamide, and case reports of a maculopapular rash and acute generalized exanthematous pustulosis related to enzalutamide.^{4,5}

Conclusions

We have presented 2 cases of toxic skin eruptions related to apalutamide, a drug that was recently introduced into our clinical practice; our report shows correlations between histology and symptoms. This drug is expected to be more widely used in the future, and dermatologists should therefore become aware of the characteristics of adverse reactions to it in order to be able to manage both diagnosis and treatment.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

References

1. Erleada[®]. Ficha técnica del medicamento. Centro de información online de medicamentos de la Agencia Europea del Medicamento (EMA). Available from: <http://www.ema.europa.eu> [cited 2021 Aug 4].
2. Uemura H, Koroki Y, Iwaki Y, Imanaka K, Kambara T, Lopez-Gitlitz A, et al. Skin rash following Administration of Apalutamide in Japanese patients with advanced prostate cancer: an integrated

- analysis of the phase 3 SPARTAN and TITAN studies and a phase 1 open-label study. *BMC Urol.* 2020;20:139.
3. Sagawa N, Watanabe Y, Mizuno Y, Takanashi S, Watanabe T, Ikeda N, et al. A case of toxic epidermal necrolysis associated with apalutamide administration. *J Cutan Immunol Allergy.* 2020;3:134–5.
 4. Saito-Sasaki N, Sawada Y, Okada E, Nakamura M. Drug eruption caused by enzalutamide: a case and literature review of androgen receptor inhibitor-related drug eruptions. *Aust J Dermatol.* 2018;59:e133–4.
 5. Alberto C, Konstantinou MP, Martinage C, Casassa E, Tournier E, Bagheri H, et al. Enzalutamide induced acute generalized exanthematous pustulosis. *J Dermatol Case Rep.* 2016;10:35–8.