



ACADEMIA ESPAÑOLA
DE DERMATOLOGÍA
Y VENEREOLOGÍA

ACTAS Dermo-Sifiliográficas

Full English text available at
www.actasdermo.org



ORIGINAL ARTICLE

Contact Allergy to Shellac. Retrospective Cross-Sectional Study With Data From the Spanish Registry of Research in Contact Dermatitis and Cutaneous Allergy (REIDAC)



P. Mercader-García^{a,*}, I. Ruiz-Gonzalez^b, R. Gonzalez-Perez^c, T. Sanz-Sanchez^d, J. Sanchez-Pérez^e, L. Borrego^f, on behalf of the REIDAC

^a Hospital General Universitario Morales Meseguer (Murcia), Universidad de Murcia, Spain

^b Complejo Hospitalario Universitario de León, Spain

^c Hospital Universitario de Araba, Spain

^d Hospital Universitario Infanta Sofía, San Sebastián de los Reyes, Madrid, Spain

^e Hospital General Universitario La Princesa, Madrid, Spain

^f Complejo Hospitalario Universitario Insular Materno Infantil, Universidad de las Palmas de Gran Canaria, Spain

Received 11 February 2023; accepted 20 February 2023

Available online 23 February 2023

KEYWORDS

Shellac;
Patch test;
Allergic contact
dermatitis;
Cross-sectional study

Abstract

Introduction: Shellac is a known allergen present mainly in cosmetics used on the eyelids and lips, although new sources of exposure have recently been described. Our objective was to assess the use of shellac as a contact allergen in Spain and the clinical profile of patients allergic to shellac.

Methods: This retrospective cross-sectional study included patients patch tested for shellac between 2018 and 2021 from the Spanish Registry of Contact Dermatitis and Cutaneous Allergy (REIDAC).

Results: A total of 980 patients were patch tested for shellac (20% in ethanol), and 37 (3.77%, 95% confidence interval [CI], 2.58–3.97%) showed positive results. Most of these patients were tested for shellac due to a suspicion of cosmetic contact dermatitis. Seven patients with present relevance were found, five with relation to cosmetics, and the other two with an occupational background of food handling. The reaction index for shellac was 0.51 and the positivity ratio was 67.56% (95% CI, 52.48–82.65%).

Conclusions: Shellac appears to be a prevalent allergen in patients with suspected contact dermatitis related with cosmetics or foodstuff. However, further studies are needed to validate its use in other patients.

© 2023 AEDV. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: pedro.mercader1@um.es (P. Mercader-García).

<https://doi.org/10.1016/j.ad.2023.02.004>

0001-7310/© 2023 AEDV. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Shellac;
Pruebas epicutáneas;
Dermatitis alérgica
de contacto;
Estudio transversal

Alergia de contacto a shellac. Estudio transversal retrospectivo con datos del Registro Español de Investigación en Dermatitis de Contacto y Alergia Cutánea (REIDAC)

Resumen

Introducción: El shellac es un alérgeno conocido por su presencia en cosméticos para labios y párpados, aunque en los últimos años se han descrito nuevas fuentes de exposición. El objetivo de nuestro trabajo fue evaluar cómo se está usando el shellac, como alérgeno de contacto en España, y las características clínicas de los pacientes alérgicos al shellac.

Métodos: Se realizó un estudio retrospectivo transversal con los pacientes incluidos en el Registro Español de Dermatitis de Contacto y Alergia Cutánea en los que se realizaron pruebas epicutáneas con shellac desde 2018 a 2021.

Resultados: El shellac (20% en etanol) fue usado en 980 pacientes, con resultados positivos en 37 de ellos (3,77%; intervalo de confianza [IC] del 95%: 2,58-3,97%). La mayoría de estos pacientes realizaron las pruebas epicutáneas por una sospecha de dermatitis de contacto por cosméticos. Se encontraron 7 pacientes con una relevancia presente, 5 de ellos en relación con el uso de cosméticos, y los otros 2 fueron dermatitis de contacto ocupacionales en el contexto de la manipulación de alimentos. El índice de reacción para el shellac fue del 0,51% y la ratio de positividad del 67,56% (IC 95%: 52,48-82,65%).

Conclusiones: El shellac parece un alérgeno frecuente en los pacientes con sospecha de dermatitis de contacto por cosméticos o alimentos. Sin embargo, se necesitan más estudios para validar su uso en otros pacientes.

© 2023 AEDV. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Shellac (CAS no. 9000-59-3, or CAS no. 97 766-50-2 for shellac wax), also known as lac, gomme laque, and lacca, is a purified resin secreted by the female of a South-east Asian insect known as *Laccifer* or *Tachardia Lacca*,¹ which has multiple uses in the cosmetic, pharmaceutical, and food industries.² In the 1990s and the first decade of this century, several cases of contact dermatitis associated with shellac use in lipsticks, eyeliners, and mascaras were described.³⁻⁷ From 2009 to 2012,^{8,9} shellac was included in the North American Contact Dermatitis Group (NACDG) baseline series, with a relatively high prevalence of positive tests; 1.6–1.7% of all patients tested, however, it was removed from the NACDG baseline series because more than 50% of positive tests were of unknown relevance and frequent irritant reactions were observed.⁹ Despite this, in 2020 shellac was included in the American Contact Dermatitis Core Allergen series¹⁰ to increase the yield of relevant positive reactions for their patients, however in most countries shellac is tested within specific series, mainly cosmetic series, when clinically suspected. Recently, several clinical reports described new sources of shellac exposure, such as tattoo ink,¹¹ a mouth guard,¹² ecological hairspray,¹³ and cases related to occupational exposure in food handlers¹⁴ due to the use of shellac as a coating agent for fruits and sweets. These cases suggest that shellac use could be more widespread than it seems.

Due to scarce comprehensive epidemiological studies outside the United States, we designed a study to investigate how shellac is being used in patch testing in Spain, what the main features of patients allergic to shellac are, relevance of patch testing, and the sources of exposure. Furthermore, shellac has been described as an irritant; therefore, another

objective was to assess the validity of patch testing with this allergen.

Material and methods

The Spanish Registry of Research in Contact Dermatitis and Cutaneous Allergy (Registro Español de Investigación en Dermatitis de Contacto y Alergia Cutánea – REIDAC) is a research project of the Spanish Group for Research in Contact Dermatitis and Skin Allergy (Grupo Español de Investigación en Dermatitis de Contacto y Alergia Cutánea – GEIDAC) which collects results of patch tests performed in 20 contact dermatitis clinics in different Spanish public hospitals; methodology and objectives of the register have been previously described.¹⁵

This cross-sectional observational study analyzed data from patients who underwent shellac patch testing from 2018 to 2021. Data were collected using OpenClinica version 3.12 (<https://www.openclinica.com/>, OpenClinica LLC, Waltham, Massachusetts, USA) and analyzed using Stata version 16 (StataCorp LLC, College Station, Texas, USA). The diagnostic suspicion in most patients was cosmetic allergic contact dermatitis, and they were studied with shellac (diluted 20% in ethanol) included in a cosmetics series supplied by Chemotechnique Diagnostics (Vellinge, Sweden) on Finn Chambers AQUA (SmartPractique, Phoenix, Arizona). In addition to the cosmetic series, all patients were studied with the Spanish baseline series, and patients with a positive patch test for fragrance mix I were also tested with a fragrance series (Chemotechnique Diagnostics, Vellinge, Sweden). The tests were performed and read according to the guidelines of the European Society of Contact Dermatitis (ESCD).¹⁶ The intensity and relevance of the positive patch tests of the allergen were assessed,

as well as the source of exposure if a present relevance was suspected. We used efficiency as an indicator of both positive test frequency and relevance, calculated as the number of tests needed to get one relevant result, and considered an allergen efficient when less than 200 tests are needed to get one relevant result.¹⁷

A descriptive analysis was performed using the χ^2 test, and data regarding MOAHLFA index (male, occupational dermatitis, atopic dermatitis, hand dermatitis, leg dermatitis, face dermatitis, age ≥ 40 years) were compared. Additionally, logistic regression with odds ratios (OR) and 95% confidence intervals were performed for the analysis of variables with significant differences between patients allergic and non-allergic to shellac. Irritant or doubtful readings were not included in MOAHLFA calculations.

To assess the validity of shellac skin tests, reaction index¹⁸ and positivity ratio¹⁹ were used. Reaction index was calculated by dividing the number of positive reactions, excluding doubtful and irritant reactions, by the number of doubtful reactions plus doubtful and irritant reactions. The result gives a number between -1 and 1 , values closer to 1 indicate that the preparation gives results that are easier to interpret and reproduce. Positivity ratio is calculated by estimating the percentage of weak positives (+) among the total number of positives (+/++/+++), which is easier to calculate than the reaction index since many weak positives are difficult to interpret and may represent irritant reactions. Ideal values for positivity ratio vary among different studies, with some authors¹⁹ stating that values above 80% indicate that the allergen is problematic, whereas others²⁰ consider values above 55% problematic.

The REIDAC study protocol was approved by the ethics committee of Complejo Hospitalario Universitario Insular Materno Infantil de las Palmas de Gran Canaria as a promoter, as well as by each involved center, and was conducted in accordance with the Declaration of Helsinki and attending local and European regulations. All patients involved in the registry signed a written consent form allowing the use of their data.

Results

During the study period, 980 patients from 15 REIDAC centers were tested with the Chemotechnique cosmetic series, with 37 patients showing positive tests to shellac. Of those with positive tests, 25 cases (67.6%) had a weak reaction (+) and 12 (32.4%) had a strong reaction (++) , the prevalence of positive test was 3.7% (95% CI, 2.5–3.9%). Additionally, 9 patients had doubtful reactions (+/–) and in the other 3 patients, reactions were qualified as irritant, the proportion of these doubtful and irritant reactions was 1.2% (95% CI, 0.5–1.9%). In 7 patients (18.9%) with positive patch tests relevance was present (Table 1), while relevance was unknown in the rest. In 2 patients, the present relevance was occupational, related to handling of lemons and sweets, while in the other 5 it was related to cosmetics use. The number of tests needed to get one positive relevant result (efficiency) was 140. No significant differences were found in the MOAHLFA index between allergic and non-allergic individuals (Table 2). The reaction index for the preparation

used was 0.51 and the positivity ratio was 67.56% (95% CI, 52.48–82.65%).

Discussion

The prevalence of shellac allergy in our series (3.7%) was higher than that reported in the NACDG,^{8,9} but similar to that recently shown by Schubert et al.²¹ in German patients, with a prevalence of 3.5% in a series of 2167 patients, although these authors included consecutive patients who underwent patch testing, while we mainly used it in patients with suspected cosmetic contact dermatitis. Other studies with patients in which the cosmetic battery was used showed higher frequencies of shellac positives than our sample; for example, the frequency was as high as 10.5%²² in a study performed by the Mayo Clinic and 11.93%²³ in a study conducted by the Hospital Universitario de Araba. We believe that the main cause of these varying prevalence rates is the different methods of patient selection for patch testing in the above-mentioned studies, which may cause different percentages of positive reactions.

In our study, we found no significant differences in any of the MOAHLFA index variables between patients allergic and non-allergic to shellac, similar to the results shown by Schubert et al.²¹; however, we found seven cases of present relevance in our series, whereas Schubert found no cases. However, in one of the patients with occupational contact dermatitis, the finding was incidental. The tests were performed due to a suspicion of contact dermatitis caused by cosmetics, when the positive test to shellac appeared we could relate it to her work in a sweet factory. The other occupational case was described in a previous article¹⁴ and was related to the use of shellac for coating lemons. In that case, there was a clinical suspicion of the source of exposure. The efficiency of patch testing was 140, suggesting that shellac is an efficient allergen at least in the subgroup of patients with suspected allergies to cosmetics or foodstuff.

Another important difference between our results and those of Schubert et al.²¹ is the high proportion of doubtful and irritant reactions found (8.5%) compared to the much lower proportion shown in our results (1.2%). Veverka et al.²² also found a low prevalence of irritant reactions (0.8%), although these authors excluded 70 patients they described as having macular erythema from their analyses, which may represent irritant reactions. Similarly, Gimenez Castillo et al.²³ did not report doubtful reactions in their study although they found that most reactions were weak. Moreover, according to Warshaw et al.,²⁰ they consider shellac as a problematic allergen because the positivity ratio in their series was 76%. However, to classify an allergen as problematic, in addition to the $\geq 55\%$ positivity ratio, Warshaw et al.²⁰ stated that the reaction index should be ≤ 0.46 . In our study, the ratio of positives was 67.56%; however, the reaction rate was 0.51, so according to the abovementioned study, shellac should be considered an indeterminate allergen. These discrepancies could be due to difficulty in interpreting positive results for this allergen.^{21,23}

Our study had several limitations, one of which is the bias in sample selection. Most of our patients underwent patch testing due to a suspected allergy to cosmetics, which

Table 1 Clinical features of patients with shellac present relevance upon patch testing.

Case number	Center	Sex	Age (years)	Main location	Occupation	Source of exposure	Other allergens
Case 1	Hospital Universitario de Araba	Female	72	Face	Retired	Cosmetics	Colophonium, 4-tert-butylphenolformaldehyde resin
Case 2	Hospital Universitario La Princesa (Madrid)	Female	37	Face	Office worker	Cosmetics	Thimerosal
Case 3	Complejo Hospitalario Universitario Las Palmas de Gran Canaria	Female	57	Face	Housewife	Cosmetics	Colophonium, nickel, fragrance mix I, cinnamyl alcohol
Case 4	Complejo Hospitalario Universitario Las Palmas de Gran Canaria	Female	74	Face	Retired	Cosmetics	Ylang ylang oil, fragrance mix I
Case 5	Complejo Hospitalario Universitario Las Palmas de Gran Canaria	Female	38	Face	Unemployed	Cosmetics	Peru balsam, budesonide, fragrance mix I, isoeugenol, ylang ylang oil
Case 6 ^a	Hospital General Universitario Morales Meseguer	Female	25	Hands	Worker in a fruit warehouse	Occupational	None
Case 7	Hospital General Universitario Morales Meseguer	Female	62	Face and hands	Worker in a sweet factory	Occupational	Tea tree oil oxidized, peppermint oil, lavender absolute, Cananga oil, Ylang Ylang oil, geranium oil, sandalwood oil, 4-tert-butylphenolformaldehyde resin, fragrance mix I

^a Patient previously published.¹³

Table 2 MOAHLFA index of patients allergic to shellac.

	Allergic, <i>n</i> (%)	Non allergic, <i>n</i> (%)	OR	95% CI
Sex, male	4 (10.81)	145 (15.37)	0.67	0.23–1.91
Occupational	6 (16.21)	77 (8.16)	2.17	0.72–5.51
Atopy	4 (10.81)	155 (16.43)	0.61	0.15–1.76
Hand	8 (21.62)	160 (16.96)	1.35	0.52–3.09
Leg	1 (2.70)	19 (2.01)	1.35	0.03–8.91
Face	17 (45.94)	509 (53.97)	0.72	0.35–1.47
Age>40 years	25 (67.56)	674 (74.20)	0.82	0.39–1.83

Statistically significant differences in qualitative variants (OR for each item) were calculated using the Chi-square test. CI, confidence interval; OR, odds ratio; MOAHLFA, male, occupational dermatitis, atopic dermatitis, hand dermatitis, leg dermatitis, face dermatitis, age \geq 40 years.

explains the presence of relevant positives, as this is the most frequently known source of exposure described in the literature. Another limitation is that the participation of several centers may lead to differences in interpretation and reading of the tests, particularly for the weak positives.²⁴ It is possible that some of the irritant reactions were not registered or appeared as weak reactions; however, it is also possible that some cases with present relevance were not

diagnosed because sources of exposure other than cosmetics were not considered. Finally, the sample size should be larger to allow better measurement of the positivity ratio and reaction index, as studies such as Warshaw et al.²⁰ excluded allergens with fewer than 100 positives from their analyses.

In conclusion, patch testing with shellac in patients with suspected cosmetic allergies, or when occupational use is

suspected, appears to be useful. Moreover, the reaction index data and positivity ratio do not clearly demonstrate that shellac is a problematic allergen, at least in these patients. However, larger studies with unbiased selected patients are required to determine the usefulness of adding this allergen to the Spanish baseline series.

Funding

REIDAC currently has funding from the Spanish Ministry of Health through an agreement with the Spanish Agency for Medicines and Health Products (AEMPS). The REIDAC accepts donations not conditioned by the pharmaceutical industry. The collaborating companies do not participate, nor have they participated, in the creation or management of the project. The collection, management, analysis, interpretation, publication and review of the data generated in the REIDAC is independent of the public and private funding sources of the project. REIDAC currently receives financial support from Sanofi.

Conflicts of interest

Pedro Mercader-García reports lectures and advisory boards from Sanofi, Leo Pharma, Lilly and Abbvie, outside the submitted work. Other authors have no conflicts of interest to disclose.

Acknowledgements

To Ignacio Garcia-Doval and Marina De Vega for their technical support.

References

- Das S, Jacob SE. Shellac. *Dermatitis*. 2011;22:220–2, <http://dx.doi.org/10.2310/6620.2011.10091>.
- Thombare N, Kumar S, Kumari U, Sakare P, Yogi RK, Prasad N, et al. Shellac as a multifunctional biopolymer: a review on properties, applications and future potential. *Int J Biol Macromol*. 2022;215:203–23, <http://dx.doi.org/10.1016/j.ijbiomac.2022.06.090>.
- Rademaker M, Kirby JD, White IR. Contact cheilitis to shellac Lanpol 5 and colophony. *Contact Dermatitis*. 1986;15:307–8, <http://dx.doi.org/10.1111/J.1600-0536.1986.TB01379.X>.
- Scheman AJ. Contact allergy to quaternium-22 and shellac in mascara. *Contact Dermatitis*. 1998;38:342, <http://dx.doi.org/10.1111/j.1600-0536.1998.tb05775.x>.
- Orton DI, Salim A, Shaw S. Allergic contact cheilitis due to shellac. *Contact Dermatitis*. 2001;44:246–63, <http://dx.doi.org/10.1034/J.1600-0536.2001.440409-5.X>.
- le Coz CJ, Leclere JM, Arnoult E, Raison-Peyron N, Pons-Guiraud A, Vigan M. Allergic contact dermatitis from shellac in mascara. *Contact Dermatitis*. 2002;46:149–52, <http://dx.doi.org/10.1034/j.1600-0536.2002.460304.x>.
- Gallo R, Marro I, Pavesi A. Allergic contact dermatitis from shellac in mascara. *Contact Dermatitis*. 2005;53:238, <http://dx.doi.org/10.1111/j.0105-1873.2005.0670f.x>.
- Warshaw EM, Belsito DV, Taylor JS, Sasseville D, DeKoven JG, Zirwas MJ, et al. North American contact dermatitis group patch test results: 2009–2010. *Dermatitis*. 2013;24:50–9, <http://dx.doi.org/10.1097/DER.0b013e318281951>.
- Warshaw EM, Maibach HI, Taylor JS, Sasseville D, DeKoven JG, Zirwas MJ, et al. North American contact dermatitis group patch test results: 2011–2012. *Dermatitis*. 2015;26:49–59, <http://dx.doi.org/10.1097/DER.000000000000097>.
- Schalock P, Dunnick C, Nedorost S, Brod B, Warshaw E, Mowad C, et al. American contact dermatitis society core allergen series: 2020 update. *Dermatitis*. 2020;31:279–82, <http://dx.doi.org/10.1097/DER.0000000000000621>.
- González-Villanueva I, Hispán Ocete P, Silvestre Salvador JF. Allergic contact dermatitis caused by a black tattoo ink in a patient allergic to shellac. *Contact Dermatitis*. 2016;75:247–8, <http://dx.doi.org/10.1111/cod.12611>.
- Melchers RC, Quint KD, van Zuuren EJ. Contact allergy to a shellac-containing mouthguard. *Contact Dermatitis*. 2022;86:554–6, <http://dx.doi.org/10.1111/COD.14081>.
- Navarro-Triviño FJ. Allergic contact dermatitis from shellac in an ecological hair spray occurring in a patient with frontal fibrosing alopecia. *Contact Dermatitis*. 2022;86:544–5, <http://dx.doi.org/10.1111/COD.14061>.
- Mercader-García P. Occupational allergic contact dermatitis caused by shellac. *Contact Dermatitis*. 2022;86:557–9, <http://dx.doi.org/10.1111/COD.14083>.
- Hernández-Fernández CP, Mercader-García P, Silvestre Salvador JF, Sánchez Pérez J, Fernández Redondo V, Miquel Miquel FJ, et al. Alérgenos candidatos para ser incluidos en la serie estándar española a partir de los datos del Registro Español de Dermatitis de Contacto. *Actas Dermosifiliogr*. 2021;112:798–805, <http://dx.doi.org/10.1016/J.AD.202105005>.
- Johansen JD, Aalto-Korte K, Agner T, Andersen KE, Bircher A, Bruze M, et al. European Society of Contact Dermatitis guideline for diagnostic patch testing – recommendations on best practice. *Contact Dermatitis*. 2015;73:195–221, <http://dx.doi.org/10.1111/cod.12432>.
- Gavín J, Mercader P, Descalzo MA, Garcia-Doval I, Silvestre JF, Sánchez-Pérez J, et al. Efficiency in patch testing: the number needed to test to get one relevant result as a new approach in the evaluation of baseline series. *Br J Dermatol*. 2020;183:391–3, <http://dx.doi.org/10.1111/bjd.18975>.
- Brasch J, Henseler T. The reaction index: a parameter to assess the quality of patch test preparations. *Contact Dermatitis*. 1992;27:203–4, <http://dx.doi.org/10.1111/J.1600-0536.1992.TB05267.X>.
- Geier J, Uter W, Lessmann H, Schnuch A. The positivity ratio – another parameter to assess the diagnostic quality of a patch test preparation. *Contact Dermatitis*. 2003;48:280–2, <http://dx.doi.org/10.1034/J.1600-0536.2003.00033.X>.
- Warshaw EM, Nelsen DD, Sasseville D, Belsito DV, Maibach HI, Zug KA, et al. Positivity ratio and reaction index: patch-test quality-control metrics applied to the North American Contact Dermatitis Group database. *Dermatitis*. 2010;21:91–7, <http://dx.doi.org/10.2310/6620.2010.09092>.
- Schubert S, Worm M, Dickel H, Wagner N, Brans R, Schröder-Kraft C, et al. Patch testing shellac in consecutive patients—data of the Information Network of Departments of Dermatology (IVDK) 2021. *Contact Dermatitis*. 2023;88:77–80, <http://dx.doi.org/10.1111/COD.14227>.
- Veverka KK, Killian JM, Yiannias JA, Hall MR, Drage LA, Davis MDP. Shellac: a tertiary care center experience. *Dermatitis*. 2018;29:226–7, <http://dx.doi.org/10.1097/DER.000000000000394>.
- Gimeno Castillo J, Rosés Gilbert P, Menéndez Parrón A, González Pérez R. Shellac: un alérgeno problemático. Experiencia durante un periodo de 5 años. *Actas Dermosifiliogr*. 2023;114:78–9, <http://dx.doi.org/10.1016/J.AD.202206017>.
- Svedman C, Isaksson M, Björk J, Mowitz M, Bruze M. 'Calibration' of our patch test reading technique is necessary. *Contact Dermatitis*. 2012;66:180–7, <http://dx.doi.org/10.1111/j.1600-0536.2011.02044.x>.