



ACTAS Dermo-Sifiliográficas

Full English text available at
www.actasdermo.org



ORIGINAL ARTICLE

Glycosaminoglycans Are Involved in the Adhesion of *Candida albicans* and *Malassezia* Species to Keratinocytes But Not to Dermal Fibroblasts[☆]



H. Ordiales,^{a,b} F. Vázquez-López,^{c,d} M. Pevida,^e B. Vázquez-Losada,^c
F. Vázquez,^{a,b,f} L.M. Quirós,^{a,b} C. Martín^{a,b,*}

^a Instituto Universitario Fernández-Vega (IUFV), Universidad de Oviedo, Oviedo, Asturias, Spain

^b Departamento de Biología Funcional, Universidad de Oviedo, Oviedo, Asturias, Spain

^c Servicio de Dermatología, Hospital Universitario Central de Asturias, Oviedo, Asturias, Spain

^d Departamento de Medicina, Universidad de Oviedo, Oviedo, Asturias, Spain

^e Centro Comunitario de Sangre y Tejidos del Principado de Asturias y CIBERER, U714, Oviedo, Asturias, Spain

^f Servicio de Microbiología, Hospital Universitario Central de Asturias, Oviedo, Asturias, Spain

Received 18 November 2020; accepted 8 February 2021

Available online 18 May 2021

KEYWORDS

Mycosis;
Glycosaminoglycans;
Infection;
Heparan sulfate;
Chondroitin sulfate

Abstract

Background and objective: Superficial mycoses are some of the most common diseases worldwide. The usual culprits—yeasts belonging to the genera *Malassezia* and *Candida*—are commensal species in the skin that can cause opportunistic infections. We aimed to determine whether these yeasts use glycosaminoglycans (GAGs) as adhesion receptors to mediate binding to epithelial cells.

Material and methods: In keratinocyte and dermal fibroblast cultures, we used rhodamine B and genistein to inhibit GAG synthesis to study the role these molecules play in the adhesion of *Candida albicans* and *Malassezia* species to cells. We also analyzed GAG involvement by means of enzyme digestion, using specific lyases.

Results: Rhodamine B partially inhibited the adhesion of both fungi to keratinocytes but not to fibroblasts. Selective digestion of heparan sulfate enhanced the binding of *Malassezia* species to keratinocytes and of both fungi to fibroblasts. Chondroitin sulfate digestion decreased *Calbicans* adhesion to keratinocytes, but increased the adhesion of the filamentous forms of this species to fibroblasts.

[☆] Please cite this article as: Ordiales H, Vázquez-López F, Pevida M, Vázquez-Losada B, Vázquez F, Quirós LM, et al. Los glicosaminoglicanos se encuentran implicados en la adherencia de *Candida albicans* y *Malassezia* spp. a queratinocitos, pero no a fibroblastos dérmicos. Actas Dermosifiliogr. 2021;112:619–624.

* Corresponding author.

E-mail address: cmartincueto@gmail.com (C. Martín).

PALABRAS CLAVE

Micosis;
Glicosaminoglicanos;
Infección;
Heparán sulfato;
Condroitín sulfato

Conclusions: Cell surface GAGs appear to play a role in the adhesion of *Calbicans* and *Malassezia* species to keratinocytes. In contrast, their adhesion to fibroblasts appears to be enhanced by GAG inhibition, suggesting that some other type of receptor is the mediator.

© 2021 Published by Elsevier España, S.L.U. on behalf of AEDV. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Los glicosaminoglicanos se encuentran implicados en la adherencia de *Candida albicans* y *Malassezia* spp. a queratinocitos, pero no a fibroblastos dérmicos

Resumen

Antecedentes y objetivo: Las micosis superficiales son algunas de las enfermedades más comunes en todo el mundo, siendo los agentes causales más frecuentes las levaduras de los géneros *Malassezia* y *Candida*, comensales habituales de la piel que pueden actuar como patógenos oportunistas. El objetivo de este trabajo es investigar si los glicosaminoglicanos (GAG) de las células epiteliales son utilizados por estos microorganismos como receptores de adhesión a las mismas.

Materiales y métodos: Se utilizaron cultivos de queratinocitos y fibroblastos dérmicos. La participación de los GAG en la adhesión de *Candida albicans* (*C. albicans*) y *Malassezia* spp. se estudió mediante inhibición específica de la síntesis de estas moléculas empleando rodamina B o genisteína. También se analizó mediante digestión enzimática in situ empleando liasas específicas.

Resultados: El tratamiento con rodamina B produjo una inhibición parcial de la adherencia de ambas especies fúngicas a queratinocitos, pero no a fibroblastos. La digestión selectiva del heparán sulfato produjo un aumento de la unión de *Malassezia* a los queratinocitos y de ambas especies a los fibroblastos. La digestión del condroitín sulfato redujo la unión de *C. albicans* en los queratinocitos, pero favoreció la unión de la forma filamentada de esta levadura en los fibroblastos.

Conclusiones: Los GAG de superficie celular de queratinocitos parecen estar implicados en la adherencia de *Candida* y *Malassezia* a la superficie celular. En los fibroblastos, por el contrario, su eliminación favorece la adherencia, sugiriendo la implicación de otro tipo de receptores.

© 2021 Publicado por Elsevier España, S.L.U. en nombre de AEDV. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The skin is colonized by a variety of microorganisms, including bacteria and, to a lesser extent, fungi, viruses, and parasites.¹ Numerous studies have described the importance of bacteria and their role in both maintaining skin homeostasis and contributing to certain types of skin infection. However, little is known about the mycobiome, the population of fungi that inhabit the body. Like bacteria, many fungi can have pathogenic effects if they proliferate excessively or the host's immune response is altered. Superficial mycoses are among the most prevalent diseases in the world, and the most commonly detected opportunistic fungi in these types of infections are yeasts of the genera *Malassezia* and *Candida*, followed by other fungi such as dermatophytes.²⁻⁴

Of the fungi found on adult skin, the most abundant belong to the genus *Malassezia*. These yeasts obtain carbon exclusively from lipids, which they are unable to synthesize, and mainly colonize sebaceous areas of skin.^{2,5} They account for more than 90% of commensal fungi present on the skin. The most abundant species are *Malassezia globosa*, *Malassezia restricta*, and *Malassezia sympodialis*. Despite their commensal nature, these species can also be associated with common skin disorders such as pityriasis versicolor and seborrheic dermatitis.^{2,5,6}

The genus *Candida* is a heterogeneous group of commensal fungi found on the skin, nails, and mucous membranes of 70% of individuals. However, in immunodeficient individuals, and in conditions conducive to immune compromise, they can become pathogens, causing superficial and even systemic infections with high mortality rates.⁷⁻⁹ Although about 20 *Candida* species are pathogens in humans, the vast majority of recorded candidiasis cases are caused by *Candida albicans*. One characteristic of members of this genus is that they are polymorphic, existing in yeast form or as hyphae or pseudohyphae.¹⁰⁻¹²

Through the skin and mucosae the human body is continuously exposed to a broad range of pathogens. To colonize these surfaces pathogens must first anchor themselves to receptors present in the host. This is a critical step that involves specific recognition between molecules on the surface of the microorganism and those that comprise the receptor.¹³ There are several cell-surface molecules that act as binding mediators, including carbohydrates, lipids, proteins, and proteoglycans (PGs).^{14,15} PGs are made up of a central protein molecule to which varying numbers of polysaccharide chains called glycosaminoglycans (GAGs) are attached. GAGs consist of varying numbers of disaccharide repeats. There are different types of GAGs, with distinct chemical structures. Two examples are heparan sulfate (HS)

and chondroitin sulfate (CS), which are formed by binding of glucuronic acid (GlcA) to N-acetylglucosamine (in the case of HS) or to N-acetylgalactosamine (in the case of CS).¹⁶ Subsequently, these chains undergo a series of post-transcriptional modifications that confer great structural diversity, enabling their involvement in numerous physiological and pathological processes.^{16,17} The interaction of GAGs with a broad range of ligands, including cytokines, growth factors, and enzymes, implicates them in processes such as cell adhesion and migration, regulation of morphogenesis, inflammation, and even cancer and various infectious processes.^{18–20} The literature describes many situations in which GAGs facilitate pathogen binding to the surface of eukaryotic cells and in some case help achieve invasion, internalization, and dissemination of the pathogen.²¹

This study investigates the role of GAGs as receptors in the process of adhesion of 2 species of commensal skin fungi, *Calbicans* and *Malassezia* species, both of which act as opportunistic pathogens in certain circumstances. The aim of the study was to determine whether GAGs are involved in the adhesion of these microorganisms to epidermal keratinocytes and dermal fibroblasts, and to examine the role in these interactions of the main GAG species present on the cell surface. The findings could help to expand current knowledge of the molecular bases of adhesion, and propose novel anti-infective strategies that block a key initial step of the pathogenic process: pathogen adhesion to target tissues. The results may have broader relevance given that adhesion is also a prerequisite for many pathogens to deliver virulence factors into host cells.

Material and Methods

Cell lines, fungal strains, and culture conditions

This research was carried out in accordance with the principles of the Declaration of Helsinki. Keratinocytes and fibroblasts were acquired from human biopsies from the tissue bank of the Community Blood and Tissue Center of Asturias after first obtaining written informed consent in accordance with Spanish laws on organ and tissue donation for research. Cells were cultured as previously indicated.²²

Calbicans and *Malassezia* species strains were obtained from clinical isolates provided by the Hospital Universitario Central de Asturias. *Calbicans* was grown at room temperature in Sabouraud (Difco) medium (BD, MD, USA) with chloramphenicol at a concentration of 50 µg/L. To obtain filamentous *Calbicans*, medium containing 0.1% glucose, 1% glycine, and 0.1% yeast extract at pH 7.5 was incubated at 37 °C and 5% CO₂ for 48 hours. *Malassezia* species were cultured at 37 °C in Sabouraud medium enriched with 1% olive oil and 1% Tween 20.

Fluorescence labeling

After overnight incubation fungi were washed 4 times with PBS. Subsequently, the cultures were resuspended in a PBS solution containing 0.1 mg/mL fluorescein isothiocyanate (FITC; Sigma Aldrich, MO, USA) at an A₆₀₀ of 0.5 and incubated in darkness for 1 hour at 37 °C. Next, the cultures were

centrifuged, washed 4 times with PBS to remove excess FITC, and resuspended in PBS at an A₆₀₀ of 0.5.

Adhesion assays

Assays to evaluate fungal adhesion to cell monolayers were carried out in 24-well plates in which cell cultures were grown to 70–90% confluence. The culture medium was removed and the cells were washed twice with PBS and blocked with 10% fetal calf serum in Dulbecco modified Eagle's minimal essential medium (DMEM) (Gibco, Life Technologies, CA, USA) for 2 hours at 37 °C and 5% CO₂. Subsequently, the wells were washed with PBS, 200 µL of the suspension containing labeled fungi was added, and the volume was made up to 500 µL with DMEM medium. The mixture was then incubated for 90 minutes at 37 °C and 5% CO₂, after which the wells were washed twice with PBS to remove unbound fungi. Finally, the cultures were disaggregated using 1% SDS and the fluorescence of the attached fungi was measured with a LS55 fluorimeter (Perkin Elmer, MA, USA), using excitation and emission wavelengths of 488 nm and 560 nm, respectively. The values from the different experiments were normalized to adhesion values obtained in the absence of any treatment, which were assigned an arbitrary value of 100.

Inhibition of GAG synthesis

Cell cultures in 24-well plates grown to approximately 70% confluence were incubated overnight in DMEM containing 50 µg/mL rhodamine B (Sigma-Aldrich) and 30 µM genistein (Sigma-Aldrich) at 37 °C. The cultures were washed twice with PBS, cell integrity was evaluated under a light microscope, and adhesion assays were carried out as described in the previous paragraph.

Enzymatic digestion of GAG on the cell surface

HS in cell cultures was digested by incubation in DMEM medium containing a mixture of heparinase I and III (500 mU/mL each; Sigma Aldrich) for 3 hours at 37 °C and 5% CO₂. CS digestion was performed in the same way, using 250 mU/mL chondroitinase ABC (Sigma Aldrich). Reactions were stopped by washing twice in PBS, cell integrity was assessed under a light microscope, and adhesion assays were then carried out as described above.

Statistical analysis

All experiments were performed at least 3 times with at least 3 replicates per group. All data were analyzed using the Statistica program (Statsoft Inc.; Tulsa, OK, USA). Differences in mean values between 2 samples were compared using the Mann-Whitney U test, with significance set at $P < 0.05$.




Figure 1 Effect of inhibition of GAG biosynthesis on yeast adhesion to skin cells. Graphs depict the effect of treatment with rhodamine B (dark bars) and genistein (light bars) on adhesion to keratinocytes (A) and fibroblasts (B). Data were normalized to yeast adhesion values recorded in untreated cells, to which an arbitrary value of 100 was assigned. Error bars represent standard deviation.

* $P < 0.05$.




Figure 2 Effect of enzymatic degradation of cellular GAG on yeast adhesion to skin cells. Graphs depict the inhibitory effect of treatment with heparinases I and III (dark bars) or chondroitinase ABC (light bars) on yeast adhesion to keratinocytes (A) and fibroblasts (B). Data were normalized to yeast adhesion values recorded in untreated cells, to which an arbitrary value of 100 was assigned. Error bars represent standard deviation.

* $P < 0.05$.

Results

Inhibition of GAG biosynthesis: effect on yeast epidermal keratinocytes and dermal fibroblasts

Treatment of keratinocytes with either rhodamine B or genistein inhibited the adhesion of both yeast and filamentous forms of *Calbicans*. However, the intensity of the effect was dependent on the inhibitor. The strongest effect was observed for rhodamine B, while genistein only reduced

adhesion values by a mean of 15% in both cases (Fig. 1A). In *Malassezia* species, rhodamine B had an effect similar to that observed for *Calbicans*, whereas treatment with genistein increased adhesion by around 17% compared with controls (Fig. 1A).

Addition of the same 2 inhibitors to dermal fibroblasts produced radically different results. Inhibition of adhesion was not observed in any case. Curiously, adhesion increased in all microbial forms analyzed. The increase in adhesion observed following genistein treatment was statistically significant (Fig. 1B).

Enzymatic digestion of GAG: effect on yeast adhesion to epidermal keratinocytes and dermal fibroblasts

Treatment of epidermal keratinocytes with heparinases I and III resulted in no significant differences in yeast adhesion with respect to controls, except in the case of *Malassezia* species, which showed an 80% increase in adhesion after elimination of HS from the surface of keratinocytes (Fig. 2A). Degradation of CS with chondroitinase ABC had the opposite effect: adhesion of *Malassezia* species was unchanged, but adhesion of both yeast and filamentous forms of *Calbicans* was significantly decreased (27% reduction in adhesion in both cases; Fig. 2A).

Digestion of HS cell on the cell surface of fibroblasts by treatment with heparinases I and III resulted in an increase in adhesion in all cases, an effect that was strongest for *Malassezia* species and was stronger for the filamentous versus the yeast form of *Calbicans* (Fig. 2A). Degradation of CS in fibroblasts resulted in no significant changes in adhesion with respect to controls, except in the case of the filamentous form of *Calbicans*, in which adhesion increased significantly (Fig. 2B).

Discussion

PGs in general and their GAG chains in particular are involved in different infectious processes, in which they act as receptors for many pathogens.²¹ The epidermis, the most superficial layer of the skin, is comprised mainly of keratinocytes, and its main function is that of a barrier that maintains homeostasis and protects against external aggressions including colonization and infection by pathogens. Immediately beneath the epidermis is the dermis, in which fibroblasts are found. This layer is less exposed to environmental factors and microorganisms. The participation of GAGs in microbial pathogenesis is conditioned by their structure,²¹ and this varies depending on the cell type and its physiological state.¹⁷ It is therefore of interest to investigate the role of these molecules as receptors in fungal infections.

To evaluate the involvement of GAGs in the interaction of *Calbicans* and *Malassezia* species with epidermal keratinocytes and dermal fibroblasts, we inhibited GAG synthesis with rhodamine B and genistein.

These inhibitors exerted distinct effects. In keratinocytes, rhodamine treatment markedly inhibited fungal adhesion. By contrast, genistein only mildly inhibited adhesion of cellular forms of *Calbicans*, and increased the adhesion of *Malassezia* species. Treatment of fibroblasts with the aforementioned inhibitors had no inhibitory effect on fungal adhesion, which was actually increased, in some cases significantly. However, the lack of significant differences was likely due to the variability of the data. The different effects of the inhibitors used are probably due to their distinct modes of action. Rhodamine B inhibits initiation and elongation of GAG chains,^{17,23,24} while genistein inhibits the activity of epidermal growth factor receptor kinase, which is required for complete expression of the genes that encode the enzymes involved in GAG production.²⁵ However, it has been reported that the effects

of genistein on GAG biosynthesis depend on GAG type and location.^{26,27} Taken together, these results show that GAGs appear to act as receptors for potentially pathogenic yeasts in keratinocytes, but not fibroblasts.

Enzymatic degradation of GAGs using different lysases showed that in keratinocytes CS appears to be more important for *Calbicans* adhesion, while elimination of HS promoted adhesion of *Malassezia* species. In dermal fibroblasts, no decrease in adhesion was observed after treatment with either inhibitor. In fact, in all cases increased adhesion was observed after elimination of HS from the cell surface. Elimination of CS by treatment with chondroitinase ABC promoted adhesion of filamentous *Calbicans*. Increased adhesion of yeast to skin cells after GAG degradation could favor greater exposure of another class of eukaryotic receptors involved in binding. On the other hand, the limited effect of degradation of individual GAG species on fungal adhesion could be indicative of cooperative phenomena involving both GAG species, as well as interaction with other receptors, as described in other epithelia.²¹

In conclusion, the present findings suggest that keratinocyte cell surface GAGs participate in *Candida* and *Malassezia* adhesion, probably in a cooperative manner. By contrast, in fibroblasts these molecules do not appear to play the same role and, in fact, interfere with adhesion, suggesting that adhesion is mediated by other receptors in these cells. The use of antifungal agents to treat these types of infections can lead to long-term resistance, underscoring the need for new therapeutic alternatives. The present findings open new avenues for the development of anti-adhesion strategies including competitive adhesion molecules that mimic GAGs, specific antibodies, and blockade of adhesin or cell receptor synthesis.

Funding

This work was funded by the 2019 'AEDV Investigates' ('AEDV investiga') prize, awarded by the Healthy Skin Foundation (Fundación Piel Sana) of the Spanish Academy of Dermatology and Venereology (AEDV).

Conflicts of Interest

The authors declare that they have no conflicts of interest.

References

1. Belkaid Y, Segre JA. Dialogue between skin microbiota and immunity. *Science*. 2014;346:954–9, <http://dx.doi.org/10.1126/science.1260144>.
2. Jo JH, Deming C, Kennedy EA, Conlan S, Polley EC, Ng WI, et al. Diverse human skin fungal communities in children converge in adulthood. *J Invest Dermatol*. 2016;136:2356–63, <http://dx.doi.org/10.1016/j.jid.2016.05.130>.
3. Silva-Rocha WP, de Azevedo MF, Chaves GM. Epidemiology and fungal species distribution of superficial mycoses in Northeast Brazil. *J Mycol Med*. 2017;27:57–64, <http://dx.doi.org/10.1016/j.mycmed.2016.08.009>. Epub 2016 Sep 20.
4. Gawdzik A, Nowogrodzka K, Hryncewicz-Gwóźdź A, Maj J, Szepietowski J, Jankowska-Konsur A. Epidemiology of dermatomycoses in southwest Poland, years

- 2011–2016. *Postepy Dermatol Alergol.* 2019;36:604–8, <http://dx.doi.org/10.5114/ada.2018.80615>.
5. Sparber F, LeibundGut-Landmann S. Host responses to *Malassezia* spp. in the mammalian skin. *Front Immunol.* 2017;8:1614, <http://dx.doi.org/10.3389/fimmu.2017.01614>, eCollection 2017.
 6. Theelen B, Cafarchia C, Gaitanis G, Bassukas ID, Boekhout T, Dawson TL Jr. *Malassezia* ecology, pathophysiology, and treatment. *Med Mycol.* 2018;56 Suppl. 1:S10–25, <http://dx.doi.org/10.1093/mmy/myx134>.
 7. Wang Y. Looking into *Candida albicans* infection, host response, and antifungal strategies. *Virulence.* 2015;6:307–8, <http://dx.doi.org/10.1080/21505594.2014.1000752>.
 8. Kashem SW, Kaplan DH. Skin immunity to *Candida albicans*. *Trends Immunol.* 2016;37:440–50, <http://dx.doi.org/10.1016/j.it.2016.04.007>.
 9. Richardson JP, Ho J, Naglik JR. *Candida*–Epithelial interactions. *J Fungi (Basel).* 2018;4:22, <http://dx.doi.org/10.3390/jof4010022>.
 10. Williams DW, Jordan RP, Wei XQ, Alves CT, Wise MP, Wilson MJ, et al. Interactions of *Candida albicans* with host epithelial surfaces. *J Oral Microbiol.* 2013;5:22434, <http://dx.doi.org/10.3402/jom.v5i0.22434>.
 11. Dadar M, Tiwari R, Karthik K, Chakraborty S, Shahali Y, Dhama K. *Candida albicans*. Biology, molecular characterization, pathogenicity, and advances in diagnosis and control—an update. *Microb Pathog.* 2018;117:128–38, <http://dx.doi.org/10.1016/j.micpath.2018.02.028>.
 12. Ribeiro FC, Rossoni RD, de Barros PP, Santos JD, Fugisaki LRO, Leão MPV, et al. Action mechanisms of probiotics on *Candida* spp. and candidiasis prevention: an update. *J Appl Microbiol.* 2020;129:175–85, <http://dx.doi.org/10.1111/jam.14511>.
 13. Pizarro-Cerdá J, Cossart P. Bacterial adhesion and entry into host cells. *Cell.* 2006;124:715–27, <http://dx.doi.org/10.1016/j.cell.2006.02.012>.
 14. Rostand KS, Esko JD. Microbial adherence to and invasion through proteoglycans. *Infect Immun.* 1997;65:1–8, <http://dx.doi.org/10.1128/IAI.65.1.1-8.1997>.
 15. Rajas O, Quirós LM, Ortega M, Vázquez-Espinosa E, Merayo-Lloves J, Vazquez F, et al. Glycosaminoglycans are involved in bacterial adherence to lung cells. *BMC Infect Dis.* 2017;17:319, <http://dx.doi.org/10.1186/s12879-017-2418-5>.
 16. Iozzo RV, Schaefer L. Proteoglycan form and function: a comprehensive nomenclature of proteoglycans. *Matrix Biol.* 2015;42:11–55, <http://dx.doi.org/10.1016/j.matbio.2015.02.003>.
 17. Esko JD, Kimata K, Lindahl U. Proteoglycans and sulfated glycosaminoglycans. In: Varki A, Cummings RD, Esko JD, Freeze HH, Stanley P, Bertozzi CR, et al., editors. *Essentials of glyco-biology.* 2nd ed. Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press; 2009 [chapter 16].
 18. Dreyfuss JL, Regatieri CV, Jarrouge TR, Cavalheiro RP, Sampaio LO, Nader HB. Heparan sulfate proteoglycans: structure, protein interactions and cell signaling. *An Acad Bras Cienc.* 2009;81:409–29, <http://dx.doi.org/10.1590/s0001-37652009000300007>.
 19. Kamhi E, Joo EJ, Dordick JS, Linhardt RJ. Glycosaminoglycans in infectious disease. *Biol Rev Camb Philos Soc.* 2013;88:928–43, <http://dx.doi.org/10.1111/brv.12034>.
 20. Fernández-Vega I, García B, García-Suárez O, Castañón S, Quirós LM. Alterations of heparan sulfate proteoglycans in cancer. *J Glycobiol.* 2015;3:1–6, <http://dx.doi.org/10.4172/2168-958X.1000109>.
 21. García B, Merayo-Lloves J, Martín C, Alcalde I, Quirós LM, Vazquez F. Surface proteoglycans as mediators in bacterial pathogens infections. *Front Microbiol.* 2016;7:220, <http://dx.doi.org/10.3389/fmicb.2016.00220>.
 22. Meana A, Iglesias J, del Rio M, Larcher F, Madrigal B, Fresno MF, et al. Large surface of cultured human epithelium obtained on a dermal matrix based on live fibroblast-containing fibrin gels. *Burns.* 1998;24:621–30, [http://dx.doi.org/10.1016/s0305-4179\(98\)00107-7](http://dx.doi.org/10.1016/s0305-4179(98)00107-7).
 23. Derrick-Roberts AL, Marais W, Byers S. Rhodamine B and 2-acetamido-1,3,6-tri-O-acetyl-4-deoxy-4-fluoro-D-glucopyranose (F-GlcNAc) inhibit chondroitin/dermatan and keratan sulphate synthesis by different mechanisms in bovine chondrocytes. *Mol Genet Metab.* 2012;106:214–20, <http://dx.doi.org/10.1016/j.ymgme.2012.04.002>.
 24. Fritz TA, Esko JD. Xyloside priming of glycosaminoglycan biosynthesis and inhibition of proteoglycan assembly. *Methods Mol Biol.* 2001;171:317–23, <http://dx.doi.org/10.1385/1-59259-209-0:317>.
 25. Piotrowska E, Jakóbkiewicz-Banecka J, Barańska S, Tyłki-Szymańska A, Czartoryska B, Wegrzyn A, et al. Genistein-mediated inhibition of glycosaminoglycan synthesis as a basis for gene expression-targeted isoflavone therapy for mucopolysaccharidoses. *Eur J Hum Genet.* 2006;14:846–52, <http://dx.doi.org/10.1038/sj.ejhg.5201623>.
 26. Nikitovic D, Tsatsakis AM, Karamanos NK, Tzanakakis GN. The effects of genistein on the synthesis and distribution of glycosaminoglycans/proteoglycans by two osteosarcoma cell lines depends on tyrosine kinase and the estrogen receptor density. *Anticancer Res.* 2003;23(1A):459–64.
 27. Martín C, Escobedo S, Suárez JE, Quirós LM. Widespread use of *Lactobacillus* OppA, a surface located protein, as an adhesin that recognises epithelial cell surface glycosaminoglycans. *Benef Microbes.* 2019;10:463–72, <http://dx.doi.org/10.3920/BM2018.0128>.